

# Circulating markers of oxidative stress and polycystic ovary syndrome (PCOS): a systematic review and meta-analysis

Mora Murri\*, Manuel Luque-Ramírez, María Insenser, Miriam Ojeda-Ojeda, and Hector F. Escobar-Morreale\*

Diabetes, Obesity and Human Reproduction Research Group, Hospital Universitario Ramón y Cajal & Universidad de Alcalá & Instituto Ramón y Cajal de Investigación Sanitaria IRYCIS & Centro de Investigación Biomédica en Red Diabetes y Enfermedades Metabólicas Asociadas (CIBERDEM), E-28034 Madrid, Spain

\*Correspondence address. moramurri@gmail.com; hescobarm.hrc@salud.madrid.org

Submitted on August 13, 2012; resubmitted on December 1, 2012; accepted on December 12, 2012

## TABLE OF CONTENTS

- Introduction
- Methods
  - Literature search
  - Inclusion and exclusion criteria
  - Data extraction
  - Quality assessment
  - Statistical analysis
- Results
  - Meta-analysis of circulating markers of oxidative stress
  - Systematic review of other oxidative stress markers
- Discussion

**BACKGROUND:** Oxidative stress might be associated with polycystic ovary syndrome (PCOS), but relatively small studies published to date do not permit reaching a definitive conclusion. We aimed at conducting a systematic review and meta-analysis of studies evaluating circulating markers of oxidative stress in patients with PCOS.

**METHODS:** We conducted a systematic review of studies reporting circulating markers of oxidative stress in women with PCOS and controls published up to June 2012, using Entrez PubMed and EMBASE online facilities. Meta-analysis calculated standardized mean differences (SMDs) and 95% confidence intervals (95CI).

**RESULTS:** From 1633 potential studies identified electronically, 68 studies, including 4933 PCOS patients and 3671 controls, were selected. For each of nine circulating markers of oxidative stress, an individual meta-analysis was conducted. Compared with control women, patients with PCOS presented higher circulating concentrations of homocysteine (23% increase, SMD 0.6, 95CI, 0.4–0.8), malondialdehyde (47% increase, SMD 1.9, 95CI 1.2–2.6) and asymmetric dimethylarginine (36% increase, SMD 1.1, 95CI 0.6–1.6), and increased superoxide dismutase activity (34% increase, SMD 1.0, 95CI 0.5–1.4) and decreased glutathione levels (50% decrease, SMD –3.7, 95CI –6.2 to –1.2) and paraoxonase-I activity (32% decrease, SMD –0.9, 95CI –1.3 to –0.4). Similar results were found when restricting the analyses to studies in which patients and controls were matched for age and body mass index.

**CONCLUSIONS:** Circulating markers of oxidative stress are abnormal in women with PCOS independent of weight excess. This finding suggests that oxidative stress may participate in the pathophysiology of this common disorder.

**Key words:** oxidative stress / polycystic ovary syndrome / hyperandrogenism / insulin resistance / obesity

## Introduction

Polycystic ovary syndrome (PCOS) is one of the most frequent endocrine and metabolic disorders, affecting 6–14% of women of child-bearing age (Knochenhauer *et al.*, 1998; Diamanti-Kandarakis *et al.*, 1999; Michelmore *et al.*, 1999; Asuncion *et al.*, 2000; March *et al.*, 2010; Sanchon *et al.*, 2012).

PCOS has been associated with metabolic disorders, including insulin resistance, obesity and diabetes. The prevalence of obesity in clinical series of patients with PCOS ranges from 30 to 75% (Ehrmann, 2005) and in as many as 28% of overweight and obese women PCOS is also present (Alvarez-Blasco *et al.*, 2006). Even in the absence of obesity, patients with PCOS frequently have excessive body fat and central adiposity (Kirchengast *et al.*, 2001) and have demonstrable insulin resistance (Goodarzi *et al.*, 2011).

PCOS and its metabolic comorbidities may be explained by the existence of a vicious circle of effects. A chronic androgen excess of ovarian and/or adrenal origin, starting prenatally or early in life, results in abdominal adiposity and android obesity in affected women (Escobar-Morreale and San Millan, 2007). Abdominal adiposity favours hypoadiponectinemia and adipose tissue dysfunction (Kadowaki and Yamauchi, 2005; Escobar-Morreale *et al.*, 2006), local and systemic cytokine excess (Gonzalez *et al.*, 1996; Fernandez-Real *et al.*, 2003) and oxidative stress (Fenkci *et al.*, 2003; Gonzalez *et al.*, 2006), among other mechanisms of disease. Abdominal adiposity thus promotes further androgen excess by a direct response of the ovaries and adrenals to inflammatory mediators or indirectly by the development of insulin resistance and compensatory hyperinsulinemia since insulin facilitates androgen secretion by these glands. Either way the vicious circle which began with abdominal adiposity is closed. Of note, the clinical heterogeneity of PCOS with respect to its metabolic complications may be explained by the individual variability in the relative contributions of androgen excess and abdominal adiposity to this vicious circle.

Aside from low-grade chronic inflammation, oxidative stress might also contribute to PCOS and its metabolic associations (Escobar-Morreale *et al.*, 2005). Oxidative stress is defined as an imbalance derived from excessive formation of oxidants in the presence of limited antioxidant defences (Turrens, 2003).

Oxidants are chemical elements that tend to gain electrons losing positive charge. They include products of normal cellular metabolism such as reactive oxygen species (ROS) and reactive nitrogen species that derive from nitric oxide (RNS). ROS derive from molecular oxygen, and include oxygen ions, free radicals (chemical species with unpaired electrons) and peroxides.

ROS and RNS are well recognized for playing a dual role, since they can be either harmful or beneficial to living systems (Valko *et al.*, 2006). At low-to-moderate concentrations, ROS and RNS are involved in physiological roles including defence against infectious agents and a number of cellular signalling systems. But when present in excess, oxidants may damage DNA, cellular lipids and proteins, interfering with their normal function (Valko *et al.*, 2007). Hence, oxidative stress participates in the pathophysiology of many human pathological conditions as well as in the physiologic process of ageing.

Protective physiological mechanisms against oxidant species involve preventative and repair mechanisms, physical defences and antioxidant defences (Valko *et al.*, 2007). Antioxidants are compounds that are

capable of disposing, scavenging or suppressing the formation of oxidants. They include enzymes [such as superoxide dismutase (SOD), catalase, glutathione peroxidase (GPx) and paraoxonase], non-enzymatic macromolecules (such as albumin and ferritin) and small molecules (such as ascorbic acid, glutathione and vitamin E).

Mounting data suggest that obesity induces oxidative stress in humans. Several studies have reported increased levels of reactive species or oxidant products in obesity (Van Gaal *et al.*, 1998; Skrha *et al.*, 1999; Dandona *et al.*, 2001; Block *et al.*, 2002; Olusi, 2002; Ozata *et al.*, 2002; Konukoglu *et al.*, 2003; Urakawa *et al.*, 2003). These levels decrease in response to weight reduction, caloric restriction and diets rich in antioxidants (Kisakol *et al.*, 2002; Uzun *et al.*, 2004; Vincent and Taylor, 2005; Murri *et al.*, 2010). Possible contributors to oxidative stress in obesity include hyperglycemia, hyperlipemia, augmented muscle activity to carry excessive weight, hyperleptinemia, chronic inflammation and inadequate antioxidant defences (Vincent and Taylor, 2005).

Furthermore, reactive species production and oxidative stress have been linked to insulin resistance. Oxidative stress impairs glucose uptake in muscle and adipose tissue, and reduces insulin secretion from pancreatic  $\beta$  cells (Matsuoka *et al.*, 1997; Rudich *et al.*, 1998; Takeda *et al.*, 2005). Recent studies have demonstrated that activation of stress-sensitive intracellular signalling pathways results in insulin resistance and impaired insulin secretion, both *in vitro* (Aguirre *et al.*, 2000; Nguyen *et al.*, 2005; Gao *et al.*, 2010) and *in vivo* (Tuncman *et al.*, 2006; Masharani *et al.*, 2011). In addition, antioxidant treatments may improve insulin sensitivity in patients with insulin resistance or type 2 diabetes (Evans and Goldfine, 2000; Evans *et al.*, 2003).

The pathogenesis of PCOS is complex and its underlying basis remains unclear. Several characteristics and associations of PCOS, including androgen excess, abdominal adiposity, insulin resistance and obesity, may contribute to the development of local and systemic oxidative stress (Vincent and Taylor, 2005; Liu *et al.*, 2010) which may reciprocally worsen these metabolic abnormalities (Fig. 1). However, previous studies addressing circulating markers of oxidative stress in women with PCOS yielded controversial results, partly because the relatively small sample sizes of most of these studies and the variability in the markers and assays used in them.


In order to overcome these limitations, we have conducted a systematic review and meta-analysis of high-quality studies addressing differences in circulating markers of oxidative stress between patients with PCOS and control women.

## Methods

### Literature search

A systematic review of the literature was conducted using Entrez PubMed and EMBASE online facilities. Studies reporting circulating markers of oxidative stress in women with PCOS and controls and published up to June 2012 were identified and analysed. In addition, a hand search of the references of the retrieved articles and relevant reviews was performed to identify other potentially eligible studies.

First, we conducted a generic search of oxidative stress and PCOS. Then, we searched for the different oxidative stress markers in blood cells, plasma and serum. The search terms were 'PCOS, polycystic ovary syndrome, oxidative stress, antioxidant, oxidant, reactive oxygen species, reactive nitrogen species, protein carbonyl, lipid peroxide,



**Figure 1** Schematic representation of mechanisms and pathways that may contribute to oxidative stress in polycystic ovary syndrome (PCOS). Several characteristics and associations of PCOS, including androgen excess, abdominal adiposity, insulin resistance and obesity, may contribute to the development of local and systemic oxidative stress which may reciprocally worsen these metabolic abnormalities. Superoxide anion can be produced by NAD(P)H oxidases, xanthine oxidase and cyclooxygenase, among others. Superoxide anion is dismutated by the SOD to hydrogen peroxide. Hydrogen peroxide is scavenged by the enzyme glutathione peroxidase which requires GSH as the electron donor, or by catalase. The oxidized glutathione is reduced back to GSH by glutathione reductase. Some transition metals can breakdown hydrogen peroxide to the reactive hydroxyl radical. The hydroxyl radical can abstract an electron from polyunsaturated fatty acid, DNA or proteins. Lipid hydroperoxides can be reduced by the reduced form of Vitamin E, which can be regenerated by Vitamin C, or by GPx using GSH as the electron donor. Nitric oxide, which is produced by nitric oxide synthase, and the superoxide anion may react together to produce peroxynitrite anion, which is a potent oxidizing agent that can cause protein, DNA and lipid oxidation. Paraonase I prevents the formation of oxidized LDL. Homocysteine might result in oxidative stress by decreasing the transcription, translation and catalytic activity of glutathione peroxidase and SOD, and by inducing inducible nitric oxide synthase. ADMA, asymmetric dimethylarginine; COX, cyclooxygenase;  $H_2O_2$ , hydrogen peroxide; HCY, homocysteine; GPx, glutathione peroxidase; GRd, glutathione reductase; GSH, reduced glutathione; GSSG, oxidized glutathione; LDL, low-density lipoprotein; NO, nitric oxide; NOS, nitric oxide synthase;  $O_2^-$  superoxide anion;  $\cdot OH$ , hydroxyl radical;  $ONOO^-$ , peroxynitrite; oxLDL, oxidized low-density lipoprotein; PON-1, paraonase I; SOD, superoxide dismutase.

malondialdehyde, 8-hydroxydeoxyguanosine, thiobarbituric acid reactive substances, nitric oxide, nitrotyrosine, sulfhydryl group, asymmetric dimethylarginine, oxidized LDL lipoprotein, xanthine oxidase, total oxidant status, total antioxidant capacity, superoxide dismutase, glutathione, glutathione peroxidase, glutathione reductase, glutathione transferase, catalase, homocystein, paraonase-I, vitamin A, vitamin C, vitamin E', including their acronyms and synonyms.

### Inclusion and exclusion criteria

Inclusion criteria included: (i) studies where reproductive-age women with PCOS were compared with non-hyperandrogenic controls for circulating markers of oxidative stress; (ii) strict diagnosis of PCOS using National Institute of Child Health and Human Development (NICHD) conference definition (Zawadzki and Dunaif, 1992), European Society of Human

Reproduction and Embryology (ESHRE)/American Society for Reproductive Medicine (ASRM) definition (The Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group, 2004), or Androgen Excess & PCOS Society (AE-PCOS) definition (Azziz et al., 2009) and (iii) articles written in English. Meta-analysis was conducted for markers studied in five or more studies. To avoid over-representation of cases, when several studies on the same series of cases and controls have been published, only the report with the larger sample size was included in the meta-analysis.

Exclusion criteria included: (i) studies in which the diagnosis of PCOS was not strict; (ii) studies addressing markers of oxidative stress in samples other than blood and its derivatives; (iii) studies reporting less than 10 cases and/or controls and (iv) articles not written in English.

Study selection was performed independently by three investigators (M.M., M.I. and M.O.-O.). Disagreements were discussed and resolved by consensus.

## Data extraction

The following data were extracted from the retrieved studies: author's names, study design, publication date, country, sample size, diagnostic criteria for PCOS, mean age and BMI, mean and standard deviation of circulating markers of oxidative stress and analytical method.

## Quality assessment

The studies included for the meta-analysis were appraised for their methodological quality according to a modification of the Newcastle-Ottawa Quality Assessment Scale for case-control studies (Wells *et al.*, 2000), as presented in Table I. The following items were evaluated in order to score the studies to maximum of eight points:

### Selection

- (1) Is the case definition adequate? (a) Yes, with independent validation [one point]; (b) yes, e.g. record linkage or based on self-reports; (c) no description.
- (2) Representativeness of the cases: (a) consecutive or obviously representative series of cases (cases with outcome of interest over a defined period of time, all cases in a defined catchment area, all cases in a defined hospital or clinic, group of hospitals, health maintenance organization, or an appropriate sample of those cases) [one point]; (b) potential for selection biases or not stated.
- (3) Selection of controls: (a) community controls [one point]; (b) hospital controls; (c) no description.
- (4) Definition of controls: (a) no history of disease (end-point) [one point]; (b) no description of source.

### Comparability

- (1) Comparability of cases and controls on the basis of design or analysis: (a) study controls for age [one point]; (b) study controls for any additional factor [one point]. Statements of no differences between groups or that the differences were not statistically significant are sufficient for establishing comparability.

### Exposure

- (1) Ascertainment of exposure: (a) secure record (PCOS diagnosis were clinically assessed) [one point]; (b) written self-report or medical record only; (c) no description.
- (2) Same method of ascertainment for cases and controls: (a) yes [one point]; (b) no.

## Statistical analysis

Meta-analyses were performed using the MIX program version 1.7 (Bax *et al.*, 2006). We used Hedge's *g* standardized mean differences (SMDs) as a measure of effect size because the outcomes of the studies were frequently measured by different assays and techniques. SMD becomes dimensionless and the scales become uniform across the different studies. Results are given as SMD and 95% confidence intervals (95CI). In order to help readers judge the practical importance of the effect found (namely, having PCOS versus being a control), we translated the effect size into Cohen's U3-50% index. This index is the percentile place difference between the two distributions (it will be 0 if there is no difference) and may be interpreted as the increase or decrease in patients compared with controls, as a percentage.

A separate random-effects model was constructed for each oxidative stress marker using the DerSimonian-Laird weighting method, which incorporates between-study variability into the calculations. This method was selected because of the considerable likelihood that there was heterogeneity and inconsistency across studies resulting from the variability in the ethnicity and race of the subjects, differences among studies in the criteria used to define PCOS, and inconsistent control of confounding factors such

as obesity. We tested heterogeneity using Cochran's *Q* and Higgins's *I*<sup>2</sup> statistics.

A subsequent subgroup analysis was restricted to studies in which the PCOS and control groups were matched for BMI and age. If a study mismatched for these variables included a subgroup of patients and controls that were matched for age and BMI, these subjects were included in sub-analysis. We asked the corresponding authors of the original articles for the mean and SD of the groups as needed (i.e. when the data were reported as medians or geometric means).

Evidence dissemination bias was estimated by funnel plot asymmetry and Egger's regression test (Egger *et al.*, 1997) when 10 or more studies were analysed. All *P* values are two-sided, and  $\alpha = 0.05$  was set as a level of statistical significance.

## Results

### Meta-analysis of circulating markers of oxidative stress

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow charts, from identification of studies to meta-analysis, are presented in Fig. 2 and the characteristics of the eligible studies for meta-analysis are summarized in Table II. The methodological quality assessment of the included studies showed that 82% of the articles were of high quality and the remaining were of medium quality (Table I).

#### Meta-analysis of homocysteine

The search yielded 487 manuscripts, of which 347 were duplicate articles, and 1 additional article was identified from the reference lists of those publications (Mohan and Vishnu, 2009). Then 59 manuscripts were immediately excluded (31 reviews, 11 congress abstracts/articles, 5 editorials, 6 letters, and 6 non-English articles). The remaining 82 studies were retrieved for a more detailed evaluation. Of these 82 studies, 33 were excluded for the following reasons: (i) not relevant for the research question (5 manuscripts); (ii) non-strict diagnosis of PCOS (Loverro *et al.*, 2002; Schachter *et al.*, 2003; Yilmaz *et al.*, 2005b; Badawy *et al.*, 2007); (iii) absence of a control group (Randeve *et al.*, 2002; Vrbikova *et al.*, 2002; Kilic-Okman and Kucuk, 2004; Kilicdag *et al.*, 2005a, b; Cagnacci *et al.*, 2006; Carlsen *et al.*, 2007; Schachter *et al.*, 2007; Gul *et al.*, 2008; Kazerooni *et al.*, 2008; Haydardedeoglu *et al.*, 2009; Makedos *et al.*, 2010; Mancini *et al.*, 2010; Palomba *et al.*, 2010; Altug Sen *et al.*, 2011; Gharakhani *et al.*, 2011; Kilic *et al.*, 2011; Rajagopal *et al.*, 2012); (iv) studies not showing the homocysteine concentrations of the control group (Luque-Ramirez *et al.*, 2009) or of the controls and patients (Glueck *et al.*, 2003); (v) homocysteine was measured in follicular fluid (Berker *et al.*, 2009) and (vi) no response to requests for data clarification (Pamuk *et al.*, 2010; Moini *et al.*, 2011, 2012). Moreover, 14 studies were excluded due to over-representation of cases also included in larger series by the same authors (Kucuk and Kilic-Okman, 2005; Yilmaz *et al.*, 2005a; Palep-Singh *et al.*, 2007, 2008; Yilmaz *et al.*, 2008; Erdogan *et al.*, 2008a; Mancini *et al.*, 2009; Kaya *et al.*, 2009a, b, c, 2010b; Arikan *et al.*, 2010; Karadeniz *et al.*, 2010, 2011).

The remaining 35 studies containing usable information were used to perform the meta-analysis of circulating homocysteine concentrations (Yarali *et al.*, 2001; Morgante *et al.*, 2002; Orio *et al.*, 2003;

**Table 1** Quality assessment of studies included in meta-analyses using a modified Newcastle–Ottawa Quality Assessment Scale for case–control studies.

Author (year)	Selection	Comparability	Exposure	Total points	Quality assessment rating <sup>a</sup>
Arikan et al. (2009)	4	2	2	8	High
Atamer et al. (2008)	3	2	2	7	High
Baskol et al. (2012)	2	2	2	6	High
Battaglia et al. (2008)	2	2	2	6	High
Bausenwein et al. (2010)	2	1	1	4	Medium
Bayrak et al. (2012)	4	2	2	8	High
Bayraktar et al. (2004)	3	2	2	7	High
Bayram et al. (2012)	2	2	2	6	High
Bickerton et al. (2005)	3	2	1	6	High
Boulman et al. (2004)	2	1	1	4	Medium
Caglar et al. (2011)	3	2	1	6	High
Cakir et al. (2011)	3	1	1	5	Medium
Carmina et al. (2005)	4	2	2	8	High
Cetinkalp et al. (2009)	2	2	1	5	Medium
Charitidou et al. (2008)	4	1	1	6	High
Demirel et al. (2007)	3	1	2	6	High
Dincer et al. (2005)	2	2	2	6	High
Dursun et al. (2006)	4	2	2	8	High
Dursun et al. (2011)	3	2	2	7	High
Espinos-Gomez et al. (2012)	4	1	2	7	High
Fan et al. (2012)	3	0	2	5	Medium
Fenkci et al. (2003)	2	2	2	6	High
Fenkci et al. (2007)	4	2	2	8	High
Fulghesu et al. (2010)	4	2	2	8	High
Guzelmeric et al. (2007)	3	2	2	7	High
Harmanci et al. (2013)	3	2	2	7	High
Hemati et al. (2011)	3	2	1	6	High
Heutling et al. (2008)	4	2	2	8	High
Karadeniz et al. (2011)	4	2	2	8	High
Karaer et al. (2010)	3	2	1	6	High
Kaya et al. (2009b)	3	2	2	7	High
Kaya et al. (2010c)	3	2	2	7	High
Kilic-Okman et al. (2004)	1	2	1	4	Medium
Kurdoglu et al. (2012)	4	2	2	8	High
Kuscu and Var, (2009)	3	2	2	7	High
Macut et al. (2006)	4	2	2	8	High
Macut et al. (2011)	2	2	2	6	High
Markou et al. (2010)	3	2	2	7	High
Mohamadin et al. (2010a)	3	2	1	6	High
Mohamadin et al. (2010b)	3	2	1	6	High
Mohan and Vishnu, (2009)	3	1	2	6	High
Moran et al. (2009)	3	2	1	6	High
Morgante et al. (2002)	2	1	2	5	Medium
Nacul et al. (2007)	3	2	2	7	High
Nafiye et al. (2010)	2	2	2	6	High
Ngo et al. (2011)	3	2	1	6	High

Continued



**Table 1** Continued

Author (year)	Selection	Comparability	Exposure	Total points	Quality assessment rating <sup>a</sup>
Oktem <i>et al.</i> (2009)	3	2	2	7	High
Orio <i>et al.</i> (2003)	2	2	2	6	High
Ozgurtas <i>et al.</i> (2008)	4	2	2	8	High
Palacio <i>et al.</i> (2006)	2	2	1	5	Medium
Rajendran <i>et al.</i> (2009)	2	1	1	4	Medium
Sabuncu <i>et al.</i> (2001)	3	2	2	7	High
Sahin <i>et al.</i> (2007)	4	2	1	7	High
Salehpour <i>et al.</i> (2011)	4	2	2	8	High
San Millan <i>et al.</i> (2006)	4	1	2	7	High
Soares <i>et al.</i> (2009)	4	2	2	8	High
Soyman <i>et al.</i> (2011)	3	2	2	7	High
Temel <i>et al.</i> (2010)	2	2	2	6	High
Topcu <i>et al.</i> (2006)	3	2	1	6	High
Torun <i>et al.</i> (2011)	1	2	1	4	Medium
Turkcuoglu <i>et al.</i> (2011)	4	1	1	6	High
Verit and Erel, (2008)	3	2	2	7	High
Victor <i>et al.</i> (2011)	3	2	2	7	High
Vrbikova <i>et al.</i> (2003)	4	0	2	6	High
Wijeyaratne <i>et al.</i> (2004)	2	1	1	4	Medium
Yarali <i>et al.</i> (2001)	2	1	2	5	Medium
Yildizhan <i>et al.</i> (2011)	4	1	1	6	High
Yilmaz <i>et al.</i> (2005c)	2	2	2	6	High

<sup>a</sup>Studies scoring 0 to 2 points were considered of low quality, from 3 to 5 points, studies were considered of medium quality, and from 6 to 8 points studies were considered of high quality.

Vrbikova *et al.*, 2003; Bayraktar *et al.*, 2004; Boulman *et al.*, 2004; Kilic-Okman *et al.*, 2004; Wijeyaratne *et al.*, 2004; Bickerton *et al.*, 2005; Carmina *et al.*, 2005; Yilmaz *et al.*, 2005c; Topcu *et al.*, 2006; Guzelmeric *et al.*, 2007; Sahin *et al.*, 2007; Atamer *et al.*, 2008; Battaglia *et al.*, 2008; Heutling *et al.*, 2008; Arikan *et al.*, 2009; Cetinkalp *et al.*, 2009; Mohan and Vishnu, 2009; Oktem *et al.*, 2009; Soares *et al.*, 2009; Fulghesu *et al.*, 2010; Karaer *et al.*, 2010; Markou *et al.*, 2010; Nafiyeh *et al.*, 2010; Temel *et al.*, 2010; Mohamadin *et al.*, 2010b; Kaya *et al.*, 2010c; Caglar *et al.*, 2011; Hemati *et al.*, 2011; Salehpour *et al.*, 2011; Bayrak *et al.*, 2012; Espinos-Gomez *et al.*, 2012; Harmanci *et al.*, 2013).

This meta-analysis included 3511 women (2090 women with PCOS and 1421 controls). Mean homocysteine concentrations were increased by 23% in women with PCOS compared with controls (SMD 0.6, 95CI 0.4–0.8,  $z = 6.5$ ,  $P < 0.01$ , Fig. 3). As suspected, heterogeneity was present ( $Q 207$ ,  $P < 0.01$ ;  $I^2 84\%$ , 95CI 78–88%), but dissemination bias was not present (Egger's regression intercept 2.3, 95CI  $-0.2$  to 4.9,  $P = 0.07$ , Fig. 3).


To rule out any influence of obesity and age in these results, we restricted the meta-analysis to studies in which patients with PCOS and controls were matched for age and BMI. We also included subgroups of patients and controls that were matched for age and BMI within studies that, overall, were not matched for age and BMI. The meta-analysis excluding six mismatched studies (Morgante *et al.*, 2002; Vrbikova *et al.*, 2003; Boulman *et al.*, 2004; Wijeyaratne *et al.*,

2004; Mohan and Vishnu, 2009; Hemati *et al.*, 2011) and maintaining only the subgroup of patients and controls presenting with similar BMI and age of another mismatched study (Espinos-Gomez *et al.*, 2012) yielded similar results (24% increase in PCOS, SMD 0.7, 95CI 0.4–0.9,  $z = 6.2$ ,  $P < 0.01$ ;  $Q 192$ ,  $P < 0.01$ ;  $I^2 85\%$ , 95CI 79–89%; Egger's regression intercept 2.4, 95CI  $-0.6$  to 5.4,  $P = 0.11$ ).

#### Meta-analysis of malondialdehyde

The initial search returned 200 manuscripts, of which 158 were duplicate articles. One additional article was identified from the reference lists of those publications (Mohan and Vishnu, 2009). Then 9 manuscripts were immediately excluded (4 congress abstracts/articles and 5 non-English articles). The remaining 34 studies were retrieved for detailed evaluation. Of these 34 studies, 18 were excluded for the following reasons: (i) not relevant for the research question (12 manuscripts); (ii) non-strict diagnosis of PCOS (Dincer *et al.*, 2001); (iii) absence of a control group (Kaya *et al.*, 2010d); (iv) malondialdehyde (MDA) was measured in follicular fluid (Yildirim *et al.*, 2007; Berker *et al.*, 2009) and (v) no response to requests for data clarification (Kaya *et al.*, 2010b; Kulkarni *et al.*, 2011). Moreover, two studies were excluded due to over-representation of cases also included in larger series by the same authors (Karadeniz *et al.*, 2008; Erdogan *et al.*, 2008b).

The remaining 14 studies containing usable information were used to perform the meta-analysis of circulating MDA concentrations



**Figure 2** Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) flowcharts of the circulating markers of oxidative stress submitted to meta-analysis. OE, oxidative stress; PCOS, polycystic ovary syndrome.

**Table II** Characteristics of the studies included in meta-analyses and in the descriptive review.

Author (year)	Country	Simple size (n)		Mean age (years)		Mean BMI (kg/m <sup>2</sup> )		Participant selection	Definition of PCOS
		PCOS	Controls	PCOS	Controls	PCOS	Controls		
<i>Arikan et al. (2009)</i>	Turkey	39	30	22.8	24.6	21.5	20.9	Age and BMI matched	ESHRE/ASRM
<i>Atamer et al. (2008)</i>	Turkey	35	30	27.3	27.9	28.0	27.6	Age and BMI matched	ESHRE/ASRM & NICHD & AE-PCOS
<i>Baskol et al. (2012)</i>	Turkey	30	20	28.2	29.5	26.1	25.1	Age and BMI matched	ESHRE/ASRM
<i>Battaglia et al. (2008)</i>	Italy	28	15	25.2	24.9	25.2	25.9	Age and BMI matched	NICHD
<i>Bausenwein et al. (2010)</i>	Germany	62	22	29.6	31.9	27.6	25.0	BMI matched	ESHRE/ASRM & NICHD & AE-PCOS
<i>Bayrak et al. (2012)</i>	Turkey	77	25	21.8	22.5	23.9	22.1	Age and BMI matched	ESHRE/ASRM
<i>Bayraktar et al. (2004)</i>	Turkey	50	25	21.4	21.6	23.9	22.1	Age and BMI matched	NICHD
<i>Bayram et al. (2012)</i>	Turkey	45	17	24.0	25.7	23.8	21.4	Age matched, no difference in BMI	ESHRE/ASRM
<i>Bickerton et al. (2005)</i>	United Kingdom	11	12	33.5	30.7	35.3	31.0	No differences in age or in BMI	NICHD
<i>Boulman et al. (2004)</i>	Israel	33	16	NA	NA	NA	NA	No differences in BMI	NICHD
<i>Caglar et al. (2011)</i>	Turkey	61	21	22.6	23.4	22.7	21.1	No differences in age or in BMI	ESHRE/ASRM
<i>Cakir et al. (2011)</i>	Turkey	52	36	23.6	25.5	24.2	22.6	Age matched, no difference in BMI	ESHRE/ASRM
<i>Carmina et al. (2005)</i>	Italy	254	127	23.7	24.1	26.4	24.3	Age matched	ESHRE/ASRM
<i>Cetinkalp et al. (2009)</i>	Turkey	129	91	24.6	25.5	24.5	24.2	Age and BMI matched	ESHRE/ASRM
<i>Charitidou et al. (2008)</i>	Greece	106	30	21.2	29.1	24.0	25.5	BMI matched	ESHRE/ASRM
<i>Demirel et al. (2007)</i>	Turkey	44	31	15.3	15.5	25.8	21.1	Age matched, only one subgroup of patients was matched for BMI	NICHD
<i>Dincer et al. (2005)</i>	Turkey	35	24	23.0	22.0	25.6	24.9	Age and BMI matched	NICHD
<i>Dursun et al. (2006)</i>	Turkey	23	23	24.4	25.1	23.0	22.2	Age and BMI matched	ESHRE/ASRM
<i>Dursun et al. (2011)</i>	Turkey	25	27	22.7	24.2	22.4	20.6	Age and BMI matched	ESHRE/ASRM & NICHD & AE-PCOS
<i>Espinos-Gomez et al. (2012)</i>	Spain	196	21	26.2	28.5	29.8	21.6	No difference in age, only one subgroup of patients showed no differences in BMI with the controls	NICHD
<i>Fan et al. (2012)</i>	China	291	281	25.0	28.4	22.9	20.9	Not matched	ESHRE/ASRM
<i>Fenkci et al. (2003)</i>	Turkey	30	31	25.8	26.1	24.3	25.2	Age matched, no difference in BMI	NICHD
<i>Fenkci et al. (2007)</i>	Turkey	31	33	24.0	25.4	24.5	23.4	Age and BMI matched	ESHRE/ASRM & NICHD & AE-PCOS
<i>Fulghesu et al. (2010)</i>	Italy	71	94	18.6	18.1	24.0	22.6	Age and BMI matched	ESHRE/ASRM
<i>Glintborg et al. (2008)</i>	Denmark	30	14	NA	NA	33.1	33.2	Age and BMI matched	NICHD
<i>Gonzalez et al. (2006)</i>	USA	16	15	27.5	31.5	29.5	31.5	Weight matched, no difference in age	ESHRE/ASRM
<i>Guzelmeric et al. (2007)</i>	Turkey	44	26	23.5	25.9	26.5	24.1	Age and BMI matched	NICHD
<i>Hamurcu et al. (2010)</i>	Turkey	36	29	21.8	23.4	26.0	23.6	Age and BMI matched	ESHRE/ASRM
<i>Harmanci et al. (2013)</i>	Turkey	23	23	22.0	21.7	21.9	21.8	Age and BMI matched	ESHRE/ASRM & NICHD & AE-PCOS
<i>Hemati et al. (2011)</i>	Iran	64	50	31.1	29.1	22.7	22.1	Age and physical activity matched	ESHRE/ASRM

Continued



Table II Continued

Author (year)	Country	Simple size (n)		Mean age (years)		Mean BMI (kg/m <sup>2</sup> )		Participant selection	Definition of PCOS
		PCOS	Controls	PCOS	Controls	PCOS	Controls		
Heutling et al. (2008)	United Kingdom	83	39	27.8	27.8	30.4	29.1	Age and BMI matched	ESHRE/ASRM
Karadeniz et al. (2011)	Turkey	98	93	24.4	25.6	24.8	24.3	Age and BMI matched	ESHRE/ASRM
Karaer et al. (2010)	Turkey	31	31	27.9	27.9	28.9	28.9	Age and BMI matched	ESHRE/ASRM
Kaya et al. (2009b)	Turkey	35	30	NA	NA	NA	NA	Age and BMI matched	ESHRE/ASRM
Kaya et al. (2010c)	Turkey	68	68	24.2	24.2	23.4	24.1	Age and BMI matched	ESHRE/ASRM
Kilic-Okman et al. (2004)	Turkey	29	25	23.9	25.2	25.3	23.3	No difference in age or BMI	NICHD
Kurdoglu et al. (2012)	Turkey	31	29	24.3	26.8	21.9	22.7	Age matched, no difference in BMI	ESHRE/ASRM
Kuscu and Var, (2009)	Turkey	31	23	23.8	22.5	21.8	20.5	Age and BMI matched	ESHRE/ASRM
Macut et al. (2006)	Serbia	179	56	24.3	24.4	25.3	23.5	Age and BMI matched	ESHRE/ASRM & NICHD & AE-PCOS
Macut et al. (2011)	Serbia	34	23	23.8	25.3	23.0	21.1	Age and BMI matched	ESHRE/ASRM
Markou et al. (2010)	Greece	17	17	25.1	25.9	20.9	21.1	Age and BMI matched	ESHRE/ASRM
Mohamadin et al. (2010a)	Saudi Arabia	50	40	30.2	29.3	29.3	25.7	No difference in age or BMI	ESHRE/ASRM
Mohamadin et al. (2010b)	Saudi Arabia	35	30	29.2	28.5	28.1	24.9	No difference in age or BMI	ESHRE/ASRM
Mohan et al. (2009)	India	56	56	NA	NA	24.8	21.1	Age matched	ESHRE/ASRM & NICHD & AE-PCOS
Moran et al. (2009)	Australia	80	27	34.1	33.8	36.0	37.4	Age and BMI matched	ESHRE/ASRM & NICHD & AE-PCOS
Morgante et al. (2002)	Italy	20	12	24.0	23.0	25.1	21.6	No difference in age	NICHD
Nacul et al. (2007)	Brazil	31	21	22.4	26.7	NA	NA	Age matched	NICHD
Nafiye et al. (2010)	Turkey	36	61	29.6	29.2	26.3	24.7	No difference in age or BMI	ESHRE/ASRM
Ngo et al. (2011)	Australia	27	20	31.5	33.1	26.9	26.3	Age and weight matched	ESHRE/ASRM
Oktem et al. (2009)	Turkey	31	31	25.6	25.6	29.7	29.3	Age and BMI matched	ESHRE/ASRM
Orio et al. (2003)	Italy	70	70	22.5	21.9	24.1	23.8	Age and BMI matched	NICHD
Ozgurtas et al. (2008)	Turkey	44	22	NA	NA	21.8	21.4	Age and BMI matched	ESHRE/ASRM
Palacio et al. (2006)	Spain	10	21	28.6	31.1	22.1	24.6	No difference in age or BMI	NICHD
Rajendran et al. (2009)	Australia	24	12	32.1	30.5	27.5	21.2	Age matched, only one subgroup of patients showed no differences in BMI with the controls	ESHRE/ASRM
Sabuncu et al. (2001)	Turkey	27	18	26.7	28.8	31.4	30.0	Age and BMI matched	NICHD
Sahin et al. (2007)	Turkey	20	20	21.6	22.9	22.4	23.2	Age and BMI matched	ESHRE/ASRM
Salehpour et al. (2011)	Iran	85	83	29.0	27.3	26.7	25.8	BMI matched	NICHD
San Millan et al. (2006)	Spain	107	58	25.0	31.0	29.6	29.5	BMI matched	NICHD
Soares et al. (2009)	Brazil	40	50	24.5	24.5	22.7	23.1	Age and BMI matched	ESHRE/ASRM
Sova et al. (2010)	Finland	50	20	27.0	30.6	27.6	26.8	Age and BMI matched	ESHRE/ASRM
Soyman et al. (2011)	Turkey	30	30	24.3	23.3	23.5	22.6	Age and BMI matched	ESHRE/ASRM
Temel et al. (2010)	Turkey	30	30	23.1	23.3	22.3	20.8	Age and BMI matched	ESHRE/ASRM
Topcu et al. (2006)	Turkey	28	26	27.1	28.8	26.6	24.7	Age matched	ESHRE/ASRM & NICHD & AE-PCOS
Torun et al. (2011)	Turkey	30	20	23.5	25.9	26.8	26.4	Age and BMI matched	ESHRE/ASRM

Turkcuoglu et al. (2011)	Turkey	40	35	25.7	27.9	24.9	23.4	No difference in BMI	ESHRE/ASRM
Verit and Erel, (2008)	Turkey	63	58	24.4	24.8	21.2	21.8	Age and BMI matched	ESHRE/ASRM
Victor et al. (2011)	Spain	39	43	24.2	26.0	22.2	21.8	Age and BMI matched	ESHRE/ASRM
Vrbikova et al. (2003)	Czech Republic	40	11	25.8	33.0	27.8	22.3	Not matched	NICHD
Wijayarathne et al. (2004)	United Kingdom	161	78	27.6	32.7	29.0	23.7	Not matched	NICHD
Yarali et al. (2001)	Turkey	30	30	27.9	31.4	27.3	25.0	BMI matched	NICHD
Yildizhan et al. (2011)	Turkey	57	27	26.4	25.4	26.9	23.9	Age matched, only one subgroup of patients showed no differences in BMI with the controls	ESHRE/ASRM
Yilmaz et al. (2005c)	Turkey	50	35	24.1	24.4	21.9	22.5	Age and weight matched	ESHRE/ASRM

AE-PCOS, Androgen Excess & Polycystic Ovary Syndrome Society; BMI, body mass index; ESHRE/ASRM, European Society of Human Reproduction and Embryology/American Society for Reproductive Medicine; NA, not available; NICHD, National Institute of Child Health and Human Development; PCOS, polycystic ovary syndrome.

(Sabuncu et al., 2001; Yilmaz et al., 2005c; Dursun et al., 2006; Palacio et al., 2006; Kusu and Var, 2009; Mohan and Vishnu, 2009; Rajendran et al., 2009; Kaya et al., 2009b; Mohamadin et al., 2010a; Karadeniz et al., 2011; Macut et al., 2011; Bayram et al., 2012; Fan et al., 2012; Kurdoglu et al., 2012). The meta-analysis included 1481 women (790 women with PCOS and 691 controls). Mean MDA levels were increased by 47% in women with PCOS compared with controls (SMD 1.9, 95CI 1.2–2.6,  $z = 5.2$ ,  $P < 0.01$ , Fig. 4), yet there was evidence of dissemination bias for this marker (Egger's regression intercept 6.7, 95CI 2.4–11.1,  $P < 0.01$ , Fig. 4). In addition, high heterogeneity was observed ( $Q = 392$ ,  $P < 0.01$ ;  $I^2 = 97%$ , 95CI 96–98%).

The meta-analysis excluding two studies mismatched for age or BMI (Mohan and Vishnu, 2009; Fan et al., 2012) and maintaining only the subgroup of patients and controls presenting with similar BMI and age of another mismatched study (Rajendran et al., 2009) yielded similar results (43% relative increase in PCOS, SMD 1.5, 95CI 0.8–2.2,  $z = 4.0$ ,  $P < 0.01$ ;  $Q = 210$ ,  $P < 0.01$ ;  $I^2 = 95%$ , 95CI 92–96%; Egger's regression intercept 9.7, 95CI 4.5–14.9,  $P < 0.01$ ).

#### Meta-analysis of asymmetric dimethylarginine (ADMA)


The initial search yielded 193 manuscripts, of which 154 were duplicate articles, and 17 manuscripts were immediately excluded (4 reviews, 8 congress abstracts/articles, 3 letters and 2 editorials). The remaining 22 studies were retrieved for a more detailed evaluation. Of these 22 studies, 9 were excluded for the following reasons: (i) not relevant for the research question (4 studies); (ii) studies not showing the ADMA concentrations of the control group (Kebapcilar et al., 2009; Teede et al., 2010; Kilic et al., 2011) and (iii) no response to requests for data clarification (Pamuk et al., 2010; Lakhani et al., 2011). Furthermore, one study was excluded due to over-representation of cases (Moran et al., 2011).

The remaining 12 studies (Demirel et al., 2007; Charitidou et al., 2008; Heutling et al., 2008; Ozgurtas et al., 2008; Moran et al., 2009; Rajendran et al., 2009; Mohamadin et al., 2010b; Ngo et al., 2011; Soyman et al., 2011; Turkcuoglu et al., 2011; Yildizhan et al., 2011; Bayrak et al., 2012) containing usable information were used to perform the ADMA meta-analysis. This meta-analysis included 1000 women (662 women with PCOS and 338 controls). Mean ADMA levels were 36% higher in women with PCOS compared with controls (SMD 1.1, 95CI 0.6–1.6,  $z = 4.2$ ,  $P < 0.01$ , Fig. 4), yet there was evidence of dissemination bias for this meta-analysis (Egger's regression intercept 10.6, 95CI 1.5–19.7,  $P = 0.03$ , Fig. 4). Heterogeneity was also observed ( $Q = 132$ ,  $P < 0.01$ ;  $I^2 = 92%$ , 95CI 87–95%).

The meta-analysis excluding one mismatched study (Turkcuoglu et al., 2011) and maintaining only the subgroup of patients and controls presenting with similar BMI and age of three mismatched studies (Demirel et al., 2007; Rajendran et al., 2009; Yildizhan et al., 2011) yielded similar results but without statistically significant evidence of dissemination bias (37% increase in PCOS, SMD 1.1, 95CI 0.6–1.7,  $z = 3.9$ ,  $P < 0.01$ ;  $Q = 128$ ,  $P < 0.01$ ;  $I^2 = 92%$ , 95CI 88–95%; Egger's regression intercept 8.6, 95CI –0.3 to 17.5,  $P = 0.06$ ).

#### Meta-analysis of nitric oxide (NO)

The initial search yielded 221 manuscripts, of which 115 were duplicate articles. Then 63 manuscripts were immediately excluded (30 reviews, 21 congress abstracts/articles, 5 editorials, 4 letters



**Figure 3** Meta-analysis of studies addressing circulating homocysteine levels (in  $\mu\text{mol/l}$ ) in patients with PCOS and control women. Results are shown as Hedges' g standardized mean differences (SMDs) and Cohen's U3–50% index. Evidence of dissemination bias was assessed using funnel plot asymmetry and Egger's regression.

and 3 non-English articles). The remaining 43 studies were retrieved for a more detailed evaluation. Of the 43 studies, 33 were excluded for the following reasons: (i) not relevant for the research question (30 studies) and (ii) studies not showing nitric oxide concentrations of the control group (Battaglia et al., 2010; Kaya et al., 2010a; Kilic et al., 2011). Moreover, two studies were excluded as over-representation of cases (Karadeniz et al., 2008; Erdogan et al., 2008b).

The remaining eight studies (Nacul et al., 2007; Kuscu and Var, 2009; Mohamadin et al., 2010b; Dursun et al., 2011; Karadeniz et al., 2011; Turkcuoglu et al., 2011; Baskol et al., 2012; Bayram et al., 2012) containing usable information were used to perform a meta-analysis that included 626 women (350 women with PCOS and 276 controls). There was no statistically significant difference in NO levels in women with PCOS compared with controls (SMD  $-0.3$ , 95CI  $-0.8$  to  $0.3$ ,  $z = 1.0$ ,  $P = 0.32$ , Fig. 4). Heterogeneity was observed ( $Q = 74$ ,  $P < 0.01$ ;  $I^2 = 91\%$ , 95CI 84–94%).

The meta-analysis excluding one mismatched study (Turkcuoglu et al., 2011) yielded similar results (SMD  $-0.4$ , 95CI  $-1.0$  to  $0.2$ ,  $z = 1.3$ ,  $P = 0.21$ ;  $Q = 68$ ,  $P < 0.01$ ;  $I^2 = 91\%$ , 95CI 84–95%).


#### Meta-analysis of glutathione

The initial search yielded 183 manuscripts, of which 120 were duplicate articles. In the reference list of those publications, one additional

article was identified for the current review (Mohan and Vishnu, 2009). Then 30 manuscripts were immediately excluded (17 reviews, 8 congress abstracts/articles, 2 letters, 1 short survey and 2 non-English articles). The remaining 34 studies were retrieved for a more detailed evaluation. Of these 34 studies, 28 were excluded for the following reasons: (i) not relevant for the research question (25 studies); (ii) non-strict diagnosis of PCOS (Dincer et al., 2001); (iii) studies not showing the glutathione concentrations of the control group (Taskin et al., 1999; Dona et al., 2012). Moreover, one study was excluded to avoid over-representation of cases (Victor et al., 2009).

The remaining five studies containing usable information were used to perform the glutathione meta-analysis (Sabuncu et al., 2001; Dincer et al., 2005; Mohan and Vishnu, 2009; Victor et al., 2011; Kurdoglu et al., 2012), including 358 women (188 women with PCOS and 170 controls). The mean glutathione levels were 50% lower in women with PCOS compared with controls (SMD  $-3.7$ , 95CI  $-6.2$  to  $-1.2$ ,  $z = 2.9$ ,  $P < 0.01$ , Fig. 5). Heterogeneity was observed ( $Q = 231$ ,  $P < 0.01$ ;  $I^2 = 98\%$ , 95CI 97–99%).

The meta-analysis excluding one mismatched study (Mohan and Vishnu, 2009) yielded similar results (49% decrease in PCOS, SMD  $-2.4$ , 95CI  $-4.3$  to  $-0.5$ ,  $z = 2.4$ ,  $P < 0.01$ ;  $Q = 98.8$ ,  $P < 0.01$ ;  $I^2 = 97\%$ , 95CI 95–98%).



**Figure 4** Meta-analysis of studies addressing circulating malondialdehyde, asymmetric dimethylarginine and nitric oxide levels in patients with PCOS and control women. All units are  $\mu\text{mol/l}$  except  $^*\mu\text{mol/mol Hb}$  and  $^{\dagger}\text{nmol/g Hb}$ . Results are shown as Hedges' g SMDs and Cohen's U3–50% index. Evidence of dissemination bias was assessed by funnel plot asymmetry and Egger's regression only for malondialdehyde and asymmetric dimethylarginine, because less than 10 studies were included in the nitric oxide meta-analysis.

*Meta-analysis of GPx activity*

The initial search yielded 64 manuscripts, of which 44 were duplicate articles. One additional article (Mohan and Vishnu, 2009) was identified from the reference list of those publications. Then 10 manuscripts were immediately excluded (6 reviews, 3 congress abstracts/articles and 1 letter). Of the remaining 11 studies, 6 were excluded for the following reasons: (i) not relevant for the research question (5 studies) and (ii) one study where the GPx activity of the control group was not shown and where GPx activity was measured in peritoneal tissues (Taskin et al., 1999).


The remaining five studies (Sabuncu et al., 2001; Mohan and Vishnu, 2009; Bausenwein et al., 2010; Macut et al., 2011; Baskol et al., 2012) containing usable information were used to perform the GPx meta-analysis that included 348 women (169 women with PCOS and 179 controls). There was no statistically significant difference in GPx activity in women with PCOS compared with controls (SMD 0.5, 95CI -0.4 to 1.4,  $z = 1.2$ ,  $P = 0.24$ , Fig. 5). Heterogeneity was observed ( $Q = 55$ ,  $P < 0.01$ ;  $I^2 = 93\%$ , 95CI 86–96%). The

meta-analysis excluding two mismatched studies (Mohan and Vishnu, 2009; Bausenwein et al., 2010) yielded similar results with no heterogeneity (SMD = 0.2, 95CI -0.2 to 0.5,  $z = 1.0$ ,  $P = 0.34$ ;  $Q = 1.94$ ,  $P = 0.38$ ;  $I^2 = 0\%$ , 95CI 0–90%).

*Meta-analysis of SOD activity*

The initial search yielded 99 manuscripts, of which 71 were duplicate articles. In the reference list of those publications, one additional article was identified for the current review (Mohan and Vishnu, 2009). Then 12 manuscripts were immediately excluded (5 reviews, 4 congress abstracts/articles, 1 letter and 2 non-English articles). The remaining 17 studies were retrieved for a more detailed evaluation. Of these 17 studies, 12 were excluded for the following reasons: (i) not relevant for the research question (eleven studies) and (ii) one study not showing SOD activity of the control group and where SOD activity was measured in peritoneal tissues (Taskin et al., 1999).





**Figure 5** Meta-analysis of studies addressing circulating antioxidant molecules, including glutathione levels, glutathione peroxidase, SOD and paraoxonase-I activities, and TAC, in patients with PCOS and control women. Units are given in the figure. Results are shown as Hedges' g SMDs and Cohen's U3-50% index. SOD, superoxide dismutase activity; TAC, total antioxidant capacity.



The remaining five studies (Sabuncu *et al.*, 2001; Kuscu and Var, 2009; Mohan and Vishnu, 2009; Bausenwein *et al.*, 2010; Macut *et al.*, 2011) containing usable information were used to perform the SOD meta-analysis that included 352 women (170 women with PCOS and 182 controls). Mean SOD activity was 34% higher in women with PCOS compared with controls (SMD = 1.0, 95CI 0.5–1.4,  $z = 4.2$ ,  $P < 0.01$ , Fig. 5). Heterogeneity was observed ( $Q = 15$ ,  $P < 0.01$ ;  $I^2 = 73\%$ , 95CI 32–89%).

The meta-analysis excluding two mismatched studies (Bausenwein *et al.*, 2010; Mohan and Vishnu, 2009) yielded similar results but without heterogeneity (31% increase in PCOS, SMD = 0.9, 95CI 0.6–1.2,  $z = 5.2$ ,  $P < 0.01$ ;  $Q = 1.1$ ,  $P = 0.58$ ;  $I^2 = 0\%$ , 95CI 0–90%).

#### Meta-analysis of paraoxonase 1 (PON-1) activity

The initial search yielded 130 manuscripts, of which 105 were duplicate articles. Then 11 manuscripts were immediately excluded (5 reviews, 3 congress abstracts/articles and 3 non-English articles). The remaining 14 studies were retrieved for a more detailed evaluation. Of these, 6 studies were excluded as they were not relevant for the research question.

The remaining eight studies (Dursun *et al.*, 2006; San Millan *et al.*, 2006; Fenkci *et al.*, 2007; Mohamadin *et al.*, 2010a; Soyman *et al.*, 2011; Torun *et al.*, 2011; Baskol *et al.*, 2012; Bayram *et al.*, 2012) which contained usable information were included in the PON-1 meta-analysis that included 562 women (331 women with PCOS and 231 controls).

Mean PON-1 activity was 32% lower in women with PCOS compared with controls (SMD  $-0.9$ , 95CI  $-1.3$  to  $-0.4$ ,  $z = 3.8$ ,  $P < 0.01$ , Fig. 5). Heterogeneity was observed ( $Q = 41$ ,  $P < 0.01$ ;  $I^2 = 83\%$ , 95CI 68–91%).

The meta-analysis excluding one mismatched study (San Millan *et al.*, 2006) yielded similar results with moderate heterogeneity (34% decrease in PCOS, SMD  $-1.0$ , 95CI  $-1.4$  to  $-0.6$ ,  $z = 4.8$ ,  $P < 0.01$ ;  $Q = 21$ ,  $P < 0.01$ ;  $I^2 = 72\%$ , 95CI 39–87%).

#### Meta-analysis of total antioxidant capacity

The initial search yielded 56 manuscripts, of which 41 were duplicate articles. The remaining 15 studies were retrieved for a more detailed evaluation. Of these 15 studies, 9 were excluded for the following reasons: (i) not relevant for the research question (7 articles); (ii) studies not showing the total antioxidant capacity (TAC) concentrations of the control group (Verit *et al.*, 2007) and (iii) TAC was measured in follicular fluid (Chattopadhyay *et al.*, 2010).

The remaining six studies containing usable information were used to perform the TAC meta-analysis (Fenkci *et al.*, 2003; Yilmaz *et al.*, 2005c; Verit and Erel, 2008; Mohamadin *et al.*, 2010a; Cakir *et al.*, 2011; Torun *et al.*, 2011). This meta-analysis included 470 women (260 women with PCOS and 210 controls). There was no significant difference in TAC in women with PCOS compared with controls (SMD  $-0.7$ , 95CI  $-2.0$  to  $0.6$ ,  $z = 1.1$ ,  $P < 0.28$ , Fig. 5), and heterogeneity was observed ( $Q = 188$ ,  $P < 0.01$ ;  $I^2 = 97\%$ , 95CI 96–98%).

### Systematic review of other oxidative stress markers

Several other circulating markers that have been studied in less than five studies with usable information are reviewed below.

Increased total oxidant status in patients with PCOS compared with controls has been reported in two studies (Verit and Erel, 2008; Torun *et al.*, 2011), yet no differences were found in another (Cakir *et al.*, 2011). Also, polymorphonuclear cells from women with PCOS produced more ROS than those of controls (Victor *et al.*, 2011).

By-products of oxidative damage would be expected to be increased in patients with PCOS if oxidative stress plays a significant role in the pathogenesis of this disorder. However, the studies conducted to date have yielded relatively conflicting results.

Plasma nitrotyrosine levels, products of nitrosative damage formed most notably by the reaction of superoxide and nitric oxide radicals with protein tyrosine residues, were significantly higher in patients with PCOS compared with healthy controls (Macut *et al.*, 2011). Serum protein carbonyl levels, a marker of oxidative modification of proteins, were increased in patients with PCOS compared with controls in two studies (Fenkci *et al.*, 2003; Kurdoglu *et al.*, 2012). Similarly, patients with PCOS presented with increased circulating concentrations of oxidized low-density lipoprotein (which promote atherogenesis through foam cell formation and inflammatory responses) in three studies (Macut *et al.*, 2006; Glintborg *et al.*, 2008; Bausenwein *et al.*, 2010), yet no differences were found between patients and controls in another study (Demirel *et al.*, 2007).

Lipid peroxides are the products of chemical damage done by oxygen free radicals to the polyunsaturated fatty acids of cell membranes. Serum lipid peroxide concentrations were increased in patients with PCOS in one study (Torun *et al.*, 2011), yet plasma thiobarbituric acid-reactive substances, a byproduct of lipid peroxidation, were similar in patients with PCOS and controls in another (Gonzalez *et al.*, 2006). However, circulating 8-hydroxydeoxyguanosine concentrations, which are a commonly used marker of oxidative stress-derived DNA damage, have been found to be similar (Hamurcu *et al.*, 2010) or even decreased (Sova *et al.*, 2010) in patients with PCOS compared with controls. Similarly, the reduction–oxidation system represented by thiol levels (also known as sulphhydryl groups) have been reported as either decreased (Torun *et al.*, 2011), similar (Baskol *et al.*, 2012) or even increased (Erdogan *et al.*, 2008b) in patients with PCOS compared with controls.

The results of studies addressing enzymes that participate in the generation or scavenging of oxidants are inconclusive. Serum xanthine oxidase, an enzyme that plays an important role in the catabolism of purines in humans and generates ROS, was increased in PCOS in one study (Baskol *et al.*, 2012). The circulating levels of glutathione-S-transferase, an enzyme that inactivates toxic endogenous compounds formed as secondary metabolites during oxidative stress, has been found to be either decreased (Kurdoglu *et al.*, 2012) or increased (Mohan and Vishnu, 2009) in patients with PCOS compared with controls. Similar conflicting results have been found for catalase, an enzyme involved in the detoxification of hydrogen peroxide, as its activity has been reported to be similar (Kurdoglu *et al.*, 2012), increased (Bausenwein *et al.*, 2010) or decreased (Mohan and Vishnu, 2009) in patients with PCOS compared with controls.

Finally, several vitamins with antioxidant properties derived from their scavenging of oxidant molecules have been studied in PCOS. Serum (Kurdoglu *et al.*, 2012) and erythrocyte (Mohan and Vishnu, 2009) vitamin C concentrations are lower in patients with PCOS compared with controls, and the same applies to circulating vitamin E concentrations (Mohan and Vishnu, 2009; Kurdoglu *et al.*, 2012). Vitamin

A and  $\beta$ -carotene, on the contrary, were not different in patients with PCOS and controls (Kurdoglu et al., 2012).

## Discussion

Many studies that have addressed oxidative stress and PCOS to date have not given a definitive conclusion about their possible association (Lee et al., 2010). With a few notable exceptions (Wijeyaratne et al., 2004; Carmina et al., 2005; San Millan et al., 2006; Charitidou et al., 2008; Espinos-Gomez et al., 2012; Fan et al., 2012), these studies have suffered from small sample sizes and included less than 100 patients with PCOS. Moreover, many different circulating markers used to estimate oxidative stress and the frequent finding of conflicting results across studies addressing the same markers may explain why the question of whether or not PCOS is associated with oxidative stress still remains open.

Our present systematic review, allowing the meta-analysis of data of nine different circulating markers of oxidative stress obtained from 68 individual studies that included a total of 4933 patients with PCOS and 3671 control women, may shed some light into this unresolved issue.

Meta-analyses of these studies showed that the concentrations of several promoters and by-products of oxidative stress were significantly increased in patients with PCOS compared with control women. Homocysteine induces oxidative stress by promoting ROS production by increasing NADPH oxidase and decreasing thioredoxin (Tyagi et al., 2005), and its concentrations were increased in PCOS patients compared with control women according to our meta-analysis. A similar increase in ADMA concentrations, which also enhances ROS production (Sydow and Munzel, 2003), and in malondialdehyde, which is an end-product of lipid peroxidation and a good marker of oxidant-mediated damage and oxidative stress (Del Rio et al., 2005), were found in PCOS.

On the other hand, our meta-analyses indicated that some circulating antioxidant markers were decreased in PCOS. Glutathione concentrations were decreased in patients with PCOS compared with controls. Glutathione plays a main protective role against oxidative stress, being a cofactor of several antioxidant enzymes, detoxifying hydrogen peroxide and lipid peroxides by the catalytic action of GPx, but being able to regenerate the most important antioxidants, vitamins C and E, back to their active forms. A similar decrease in the activity of PON-1 in patients with PCOS may contribute to oxidative stress because PON-1 is an antioxidant enzyme that prevents oxidation of lipoproteins and hydrolyzes atherogenic products of oxidative lipid modification such as phospholipid peroxides and cholesterol ester hydroperoxides. Furthermore, even the increase found in patients with PCOS in the activity of SOD, a potent protective enzyme scavenging superoxide anion radical by catalyzing its dismutation to  $H_2O_2$  and  $O_2$  (Pollack and Leeuwenburgh, 1999), may be interpreted as a compensatory mechanism in response to the increased production of oxidant molecules mentioned above.

Although meta-analyses did not result in significant differences among patients with PCOS and control women in nitric oxide levels, GPx activity and TAC, overall, the decrease in glutathione and paraoxonase-I levels in parallel with an increase of oxidant molecules such as homocysteine, ADMA and MDA suggests an imbalance in the oxidative status of patients with PCOS.

However, because obesity enhances ROS production promoting oxidative stress, and PCOS and obesity are frequently associated, the possibility exists that the association of PCOS with markers of oxidative stress could be spurious, actually deriving from the association of PCOS with obesity. The fact that the abnormalities in circulating markers of oxidative stress persisted when we restricted the meta-analyses to studies and subgroups of women matched for age and BMI indicates that oxidative stress may be present in PCOS irrespective of its well-known association with weight excess, obesity and ageing. Moreover, the systematic review identified individual studies showing differences between patients with PCOS and controls in other oxidants, by-products of oxidative stress and antioxidant vitamins and compounds, further supporting the participation of oxidative stress in PCOS.

Because PCOS is a life-time heterogenous endocrinopathy, long-term management of this frequent disorder must consider all the consequences of the syndrome, including the metabolic comorbidities in which oxidative stress may play a role. For this purpose, the finding of a clinically helpful individual marker of oxidative stress in patients with PCOS would be of importance. Our present review suggests that glutathione, paraoxonase-I, homocysteine, ADMA and MDA levels might be appropriate to estimate the oxidative status of women with PCOS. However, the magnitude of the changes in markers of oxidative stress associated with PCOS was relatively small and of uncertain clinical consequences and, therefore, routine clinical measurements of these markers cannot be recommended with the evidence gathered to date.

Regarding therapeutic strategies, correction of oxidative stress by improving antioxidant defences through body fat mass reduction, pharmacological agents, exercise and/or dietary modification might have beneficial effects in PCOS, as has been demonstrated for insulin-resistant disorders (Abdel-Wahab et al., 2002; Vincent et al., 2007; Wright and Sutherland, 2008). Although experimental data suggest that strategies targeting oxidative stress might prove useful for PCOS (Masharani et al., 2010; Rzepczynska et al., 2011), the paucity of studies addressing antioxidant therapy in patients with PCOS precludes making any recommendation for routine clinical use of antioxidants in this disorder.

The strengths of our current systematic review and meta-analysis include the wide range of markers of oxidative stress evaluated, the large number of subjects included in the analyses, the exclusion of duplicate reports and the medium- to high-quality requirement for inclusion of studies in the meta-analysis as assessed by a modified Newcastle-Ottawa Quality Assessment Scale for case-control studies (Wells et al., 2000).

However, our study is not free of limitations. First, the magnitude of the changes in circulating markers of oxidative stress associated with PCOS was modest (50% or less) and therefore the results must be interpreted with caution. Processes of oxidative stress participate in the pathogenesis of disease mostly at the tissue level and may not translate entirely into its circulating markers, explaining the modest changes observed here. Secondly, two of the largest meta-analyses showed evidence for dissemination (publication) bias, suggesting that studies with positive results (namely, those showing statistically significant differences between patients and controls) were more likely to be published than studies with negative results, another reason for a careful interpretation of the results. Thirdly, most if not all the

studies conducted to date compared patients with PCOS with control women selected as to have no evidence of androgen excess and, in some cases, other metabolic disorders. This use of 'hypernormal' controls may lead to false-positive results that could have been avoided by the more appropriate recruitment of unselected control women from the general population (Hattersley and McCarthy, 2005). Fourthly, there was considerable heterogeneity in the studies included in the meta-analyses. Factors contributing to this heterogeneity include the differences in the criteria used for the definition of PCOS, differences in race and ethnicity of the subjects and the use of different assays and different samples (plasma, serum, leukocytes and erythrocytes). Furthermore, blood sampling after an adequate fasting period is a critical issue when analysing oxidative stress as fasting may modify the results of assays (Pollack and Leeuwenburgh, 1999); however, this key issue for the assessment of markers of oxidative stress was not specified with detail in most studies.

In order to overcome the impact that the heterogeneity resulting from clinical and/or methodological issues might have had on the results of the meta-analyses we chose Hedges's *g* SMD as a measure of the effect size (Durlak, 2009) and implemented random-effects models using the DerSimonian–Laird weighting method (DerSimonian and Laird, 1986). Random-effects models assume that the estimated effects in the different studies are variable but there is some type of specific distribution, using the observed effect from individual studies to estimate this distribution. Specifically, the DerSimonian–Laird weighting method estimates the magnitude of

heterogeneity and assigns a greater variability (confidence interval) to the effect size to compensate for this heterogeneity. Of note, clinical factors were possibly responsible for the heterogeneity observed in some of the meta-analyses (i.e. GPx and SOD activities) as the sub-analysis of studies and subgroups matched for age and BMI were no longer heterogeneous. In any case, the overall results of the meta-analyses were similar including or excluding mismatched groups and studies, indicating that heterogeneity was correctly compensated by the statistical approach used here.

In conclusion, as summarized in Table III, our systematic review and meta-analysis indicates that several circulating markers of oxidative stress are abnormal in women with PCOS independent of weight excess. Although modest in magnitude, these abnormalities suggest that oxidative stress may participate in the pathophysiology of this common disorder. However, in light of the evidence available at present, neither the routine measurement of markers of oxidative stress nor the use of antioxidant therapies can be recommended as yet for the clinical management of PCOS.

## Acknowledgements

The authors are very grateful to the authors of the original articles who kindly provided additional data or clarifications about their publications.

## Authors' roles

M.M. contributed to the study conception and design and data collection and analysis, and wrote the draft of the article. M.L-R. analysed the data and revised the article critically for important intellectual content. M.I. and M.O.-O. contributed to data collection and analysis. H.F.E-M. contributed to the study conception and design, analysed the data, revised the article critically for important intellectual content and wrote the final version of the manuscript. All authors approved the final version of the article and take full responsibility for the accuracy of its content.

## Funding

This work was supported by Grants PI 080944 and PII100357 from Instituto de Salud Carlos III, Spanish Ministry of Economy and Competitiveness. M.M. is the recipient of a postdoctoral grant (Programa Sara Borrell, CD11/0030) from Instituto de Salud Carlos III, Spanish Ministry of Economy and Competitiveness.

## Conflict of interest

None declared.

## References

- Abdel-Wahab YH, O'Harte FP, Mooney MH, Barnett CR, Flatt PR. Vitamin C supplementation decreases insulin glycation and improves glucose homeostasis in obese hyperglycemic (ob/ob) mice. *Metabolism* 2002;**51**:514–517.
- Aguirre V, Uchida T, Yenush L, Davis R, White MF. The c-Jun NH(2)-terminal kinase promotes insulin resistance during association with insulin receptor substrate-1 and phosphorylation of Ser(307). *J Biol Chem* 2000;**275**:9047–9054.

**Table III Summary of the meta-analysis of circulating markers of oxidative stress in polycystic ovary syndrome (PCOS).**

Marker	Action	Changes in PCOS
<i>Promoters and byproducts of oxidative stress</i>		
Homocysteine	Promotes reactive species	↑
ADMA	Promotes reactive species	↑
Malondialdehyde	End-product of lipid peroxidation	↑
Nitric oxide	Promotes reactive nitrogen species	↔
<i>Antioxidants</i>		
Glutathione	Detoxifies hydrogen peroxide and lipid peroxides, prevents protein from oxidation	↓
Paraoxonase-I	Prevents oxidation of lipoproteins by reactive species	↓
SOD	Converts superoxide anions to hydrogen peroxide and molecular oxygen	↑
Glutathione peroxidase	Detoxifies hydrogen peroxide, peroxy nitrates and lipid peroxides	↔
TAC	Prevents oxidation and detoxifies oxidants	↔

↓, decreased; ↑, increased; ↔, unchanged; ADMA, asymmetric dimethylarginine; SOD, superoxide dismutase activity TAC, total antioxidant capacity.

- Altug Sen T, Koken R, Narci A, Yilmazer M. Homocysteine and ghrelin link with polycystic ovary syndrome in relation to obesity. *J Pediatr Adolesc Gynecol* 2011; **24**:211–217.
- Alvarez-Blasco F, Botella-Carretero JL, San Millan JL, Escobar-Morreale HF. Prevalence and characteristics of the polycystic ovary syndrome in overweight and obese women. *Arch Intern Med* 2006; **166**:2081–2086.
- Arikan S, Akay H, Bahceci M, Tuzcu A, Gokalp D. The evaluation of endothelial function with flow-mediated dilatation and carotid intima media thickness in young nonobese polycystic ovary syndrome patients; existence of insulin resistance alone may not represent an adequate condition for deterioration of endothelial function. *Fertil Steril* 2009; **91**:450–455.
- Arikan S, Bahceci M, Tuzcu A, Kale E, Gokalp D. Serum resistin and adiponectin levels in young non-obese women with polycystic ovary syndrome. *Gynecol Endocrinol* 2010; **26**:161–166.
- Asuncion M, Calvo RM, San Millan JL, Sancho J, Avila S, Escobar-Morreale HF. A prospective study of the prevalence of the polycystic ovary syndrome in unselected Caucasian women from Spain. *J Clin Endocrinol Metab* 2000; **85**:2434–2438.
- Atamer A, Demir B, Bayhan G, Atamer Y, Ilhan N, Akkus Z. Serum levels of leptin and homocysteine in women with polycystic ovary syndrome and its relationship to endocrine, clinical and metabolic parameters. *J Int Med Res* 2008; **36**:96–105.
- Azziz R, Carmina E, Dewailly D, Diamanti-Kandarakis E, Escobar-Morreale HF, Futterweit W, Janssen OE, Legro RS, Norman RJ, Taylor AE et al. The androgen excess and PCOS society criteria for the polycystic ovary syndrome: the complete task force report. *Fertil Steril* 2009; **91**:456–488.
- Badawy A, State O, El Gawad S, El Aziz OA. Plasma homocysteine and polycystic ovary syndrome: the missed link. *Eur J Obstet Gynecol Reprod Biol* 2007; **131**:68–72.
- Baskol G, Aygen E, Erdem F, Caniklioglu A, Narin F, Sahin Y, Kaya T. Assessment of paraoxonase I, xanthine oxidase and glutathione peroxidase activities, nitric oxide and thiol levels in women with polycystic ovary syndrome. *Acta Obstet Gynecol Scand* 2012; **91**:326–330.
- Battaglia C, Mancini F, Cianciosi A, Busacchi P, Facchinetti F, Marchesini GR, Marzocchi R, de Alosio D. Vascular risk in young women with polycystic ovary and polycystic ovary syndrome. *Obstet Gynecol* 2008; **111**:385–395.
- Battaglia C, Mancini F, Battaglia B, Facchinetti F, Artini PG, Venturoli S. L-arginine plus drospirenone-ethinyl estradiol in the treatment of patients with PCOS: a prospective, placebo controlled, randomised, pilot study. *Gynecol Endocrinol* 2010; **26**:861–868.
- Bausenwein J, Serke H, Eberle K, Hirrlinger J, Jogschies P, Hmeidan FA, Blumenauer V, Spanel-Borowski K. Elevated levels of oxidized low-density lipoprotein and of catalase activity in follicular fluid of obese women. *Mol Hum Reprod* 2010; **16**:117–124.
- Bax L, Yu LM, Ikeda N, Tsuruta H, Moons KG. Development and validation of MIX: comprehensive free software for meta-analysis of causal research data. *BMC Med Res Methodol* 2006; **6**:50.
- Bayrak T, Dursun P, Bayrak A, Gultekin M, Kulusari A, Cakir E, Ozyurt M, Zeyneloglu HB. Paraoxonase lactonase activity (PON-HTLase), asymmetric dimethylarginine (ADMA) and platelet activating factor-acetylhydrolase (PAF-AH) activity in non-obese women with PCOS. *Gynecol Endocrinol* 2012; **28**:874–878.
- Bayraktar F, Dereli D, Ozgen AG, Yilmaz C. Plasma homocysteine levels in polycystic ovary syndrome and congenital adrenal hyperplasia. *Endocr J* 2004; **51**:601–608.
- Bayram F, Kocer D, Ozsan M, Muhtaroglu S. Evaluation of endothelial dysfunction, lipid metabolism in women with polycystic ovary syndrome: relationship of paraoxonase I activity, malondialdehyde levels, low-density lipoprotein subfractions, and endothelial dysfunction. *Gynecol Endocrinol* 2012; **28**:497–501.
- Berker B, Kaya C, Aytac R, Satioglu H. Homocysteine concentrations in follicular fluid are associated with poor oocyte and embryo qualities in polycystic ovary syndrome patients undergoing assisted reproduction. *Hum Reprod* 2009; **24**:2293–2302.
- Bickerton AS, Clark N, Meeking D, Shaw KM, Crook M, Lumb P, Turner C, Cummings MH. Cardiovascular risk in women with polycystic ovarian syndrome (PCOS). *J Clin Pathol* 2005; **58**:151–154.
- Block G, Dietrich M, Norkus EP, Morrow JD, Hudes M, Caan B. Factors associated with oxidative stress in human populations. *Am J Epidemiol* 2002; **156**:274–285.
- Boulman N, Levy Y, Leiba R, Shachar S, Linn R, Zinder O, Blumenfeld Z. Increased C-reactive protein levels in the polycystic ovary syndrome: a marker of cardiovascular disease. *J Clin Endocrinol Metab* 2004; **89**:2160–2165.
- Caglar GS, Oztas E, Karadag D, Pabuccu R, Demirtas S. Ischemia-modified albumin and cardiovascular risk markers in polycystic ovary syndrome with or without insulin resistance. *Fertil Steril* 2011; **95**:310–313.
- Cagnacci A, Tirelli A, Renzi A, Paoletti AM, Volpe A. Effects of two different oral contraceptives on homocysteine metabolism in women with polycystic ovary syndrome. *Contraception* 2006; **73**:348–351.
- Cakir E, Ozbek M, Ozkaya E, Colak N, Cakal E, Sayki M, Gungunes A, Aliyazicioglu Y, Mentese A, Delibasi T. Oxidative stress markers are not valuable markers in lean and early age of polycystic ovary syndrome patients. *J Endocrinol Invest* 2011; **34**:e178–182.
- Carlsen SM, Kjotrod S, Vanky E, Romundstad P. Homocysteine levels are unaffected by metformin treatment in both nonpregnant and pregnant women with polycystic ovary syndrome. *Acta Obstet Gynecol Scand* 2007; **86**:145–150.
- Carmina E, Chu MC, Longo RA, Rini GB, Lobo RA. Phenotypic variation in hyperandrogenic women influences the findings of abnormal metabolic and cardiovascular risk parameters. *J Clin Endocrinol Metab* 2005; **90**:2545–2549.
- Cetinkalp S, Karadeniz M, Erdogan M, Zengi A, Cetintas V, Tetik A, Eroglu Z, Kosova B, Ozgen AG, Saygili F et al. Apolipoprotein E gene polymorphism and polycystic ovary syndrome patients in Western Anatolia, Turkey. *J Assist Reprod Genet* 2009; **26**:1–6.
- Charitidou C, Farmakiotis D, Zournatzi V, Pidonia I, Pegiou T, Karamanis N, Hatzistilianou M, Katsikis I, Panidis D. The administration of estrogens, combined with anti-androgens, has beneficial effects on the hormonal features and asymmetric dimethyl-arginine levels, in women with the polycystic ovary syndrome. *Atherosclerosis* 2008; **196**:958–965.
- Chattopadhyay R, Ganesh A, Samanta J, Jana SK, Chakravarty BN, Chaudhury K. Effect of follicular fluid oxidative stress on meiotic spindle formation in infertile women with polycystic ovarian syndrome. *Gynecol Obstet Invest* 2010; **69**:197–202.
- Dandona P, Mohanty P, Ghanim H, Aljada A, Browne R, Hamouda W. The suppressive effect of dietary restriction and weight loss in the obese on the generation of reactive oxygen species by leukocytes, lipid peroxidation, and protein carbonylation. *J Clin Endocrinol Metab* 2001; **86**:355–362.
- Del Rio D, Stewart AJ, Pellegrini N. A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *Nutr Metab Cardiovasc Dis* 2005; **15**:316–328.
- Demirel F, Bideci A, Cinaz P, Camurdan MO, Biberoglu G, Yesilkaya E, Hasanoglu A. Serum leptin, oxidized low density lipoprotein and plasma asymmetric dimethylarginine levels and their relationship with dyslipidaemia in adolescent girls with polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2007; **67**:129–134.
- DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; **7**:177–188.
- Diamanti-Kandarakis E, Kouli CR, Bergiele AT, Filandra FA, Tsianateli TC, Spina GG, Zapanti ED, Bartzis MI. A survey of the polycystic ovary syndrome in the Greek island of Lesbos: hormonal and metabolic profile. *J Clin Endocrinol Metab* 1999; **84**:4006–4011.
- Dincer Y, Ozen E, Kadioglu P, Hatemi H, Akcay T. Effect of sex hormones on lipid peroxidation in women with polycystic ovary syndrome, healthy women, and men. *Endocr Res* 2001; **27**:309–316.
- Dincer Y, Akcay T, Erdem T, Ilker Saygili E, Gundogdu S. DNA damage, DNA susceptibility to oxidation and glutathione level in women with polycystic ovary syndrome. *Scand J Clin Lab Invest* 2005; **65**:721–728.
- Dona G, Sabbadin C, Fiore C, Bragadin M, Giorgino FL, Ragazzi E, Clari G, Bordin L, Armanini D. Inositol administration reduces oxidative stress in erythrocytes of patients with polycystic ovary syndrome. *Eur J Endocrinol* 2012; **166**:703–710.
- Durlak JA. How to select, calculate, and interpret effect sizes. *J Pediatr Psychol* 2009; **34**:917–928.
- Dursun P, Demirtas E, Bayrak A, Yarali H. Decreased serum paraoxonase I (PON1) activity: an additional risk factor for atherosclerotic heart disease in patients with PCOS?. *Hum Reprod* 2006; **21**:104–108.
- Dursun E, Akalin FA, Guncu GN, Cinar N, Aksoy DY, Tozum TF, Kilinc K, Yildiz BO. Periodontal disease in polycystic ovary syndrome. *Fertil Steril* 2011; **95**:320–323.
- Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *Br Med J* 1997; **315**:629–634.
- Ehrmann DA. Polycystic ovary syndrome. *N Engl J Med* 2005; **352**:1223–1236.



- Erdogan M, Karadeniz M, Alper GE, Tamsel S, Uluer H, Caglayan O, Saygili F, Yilmaz C. Thrombin-activatable fibrinolysis inhibitor and cardiovascular risk factors in polycystic ovary syndrome. *Exp Clin Endocrinol Diabetes* 2008a; **116**:143–147.
- Erdogan M, Karadeniz M, Berdeli A, Alper G, Caglayan O, Yilmaz C. The relationship of the interleukin-6-174 G>C gene polymorphism with oxidative stress markers in Turkish polycystic ovary syndrome patients. *J Endocrinol Invest* 2008b; **31**:624–629.
- Escobar-Morreale HF, Luque-Ramirez M, San Millan JL. The molecular-genetic basis of functional hyperandrogenism and the polycystic ovary syndrome. *Endocr Rev* 2005; **26**:251–282.
- Escobar-Morreale HF, Villuendas G, Botella-Carretero JJ, Alvarez-Blasco F, Sanchon R, Luque-Ramirez M, San Millan JL. Adiponectin and resistin in PCOS: a clinical, biochemical and molecular genetic study. *Hum Reprod* 2006; **21**:2257–2265.
- Escobar-Morreale HF, San Millan JL. Abdominal adiposity and the polycystic ovary syndrome. *Trends Endocrinol Metab* 2007; **18**:266–272.
- Espinos-Gomez JJ, Rodriguez-Espinosa J, Ordonez-Llanos J, Calaf-Alsina J. Metabolic syndrome in Mediterranean women with polycystic ovary syndrome: when and how to predict its onset. *Gynecol Endocrinol* 2012; **28**:264–268.
- Evans JL, Goldfine ID. Alpha-lipoic acid: a multifunctional antioxidant that improves insulin sensitivity in patients with type 2 diabetes. *Diabetes Technol Ther* 2000; **2**:401–413.
- Evans JL, Goldfine ID, Maddux BA, Grodsky GM. Are oxidative stress-activated signaling pathways mediators of insulin resistance and beta-cell dysfunction?. *Diabetes* 2003; **52**:1–8.
- Fan P, Liu H, Wang Y, Zhang F, Bai H. Apolipoprotein E-containing HDL-associated platelet-activating factor acetylhydrolase activities and malondialdehyde concentrations in patients with PCOS. *Reprod Biomed Online* 2012; **24**:197–205.
- Fenki V, Fenki S, Yilmazer M, Serteser M. Decreased total antioxidant status and increased oxidative stress in women with polycystic ovary syndrome may contribute to the risk of cardiovascular disease. *Fertil Steril* 2003; **80**:123–127.
- Fenki IV, Serteser M, Fenki S, Kose S. Paraoxonase levels in women with polycystic ovary syndrome. *J Reprod Med* 2007; **52**:879–883.
- Fernandez-Real JM, Broch M, Vendrell J, Ricart W. Insulin resistance, inflammation, and serum fatty acid composition. *Diabetes Care* 2003; **26**:1362–1368.
- Fulghesu A, Magnini R, Portoghese E, Angioni S, Minerba L, Melis GB. Obesity-related lipid profile and altered insulin secretion in adolescents with polycystic ovary syndrome. *J Adolesc Health* 2010; **46**:474–481.
- Gao D, Nong S, Huang X, Lu Y, Zhao H, Lin Y, Man Y, Wang S, Yang J, Li J. The effects of palmitate on hepatic insulin resistance are mediated by NADPH oxidase 3-derived reactive oxygen species through JNK and p38MAPK pathways. *J Biol Chem* 2010; **285**:29965–29973.
- Gharakhani M, Neghab N, Farimani M. Is reducing ovarian volume in polycystic ovarian syndrome patients after administration of metformin associated with improving cardiovascular risk factors? *Int J Fertil Steril* 2011; **5**:90–95.
- Glintborg D, Hojlund K, Andersen M, Henriksen JE, Beck-Nielsen H, Handberg A. Soluble CD36 and risk markers of insulin resistance and atherosclerosis are elevated in polycystic ovary syndrome and significantly reduced during pioglitazone treatment. *Diabetes Care* 2008; **31**:328–334.
- Glueck CJ, Wang P, Bornovali S, Goldenberg N, Sieve L. Polycystic ovary syndrome, the G1691A factor V Leiden mutation, and plasminogen activator inhibitor activity: associations with recurrent pregnancy loss. *Metabolism* 2003; **52**:1627–1632.
- Gonzalez F, Chang L, Horab T, Lobo RA. Evidence for heterogeneous etiologies of adrenal dysfunction in polycystic ovary syndrome. *Fertil Steril* 1996; **66**:354–361.
- Gonzalez F, Rote NS, Minium J, Kirwan JP. Reactive oxygen species-induced oxidative stress in the development of insulin resistance and hyperandrogenism in polycystic ovary syndrome. *J Clin Endocrinol Metab* 2006; **91**:336–340.
- Goodarzi MO, Dumesic DA, Chazenbalk G, Azziz R. Polycystic ovary syndrome: etiology, pathogenesis and diagnosis. *Nat Rev Endocrinol* 2011; **7**:219–231.
- Gul OB, Somunkiran A, Yucler O, Demirci F, Ozdemir I. The effect of ethinyl estradiol-cyproterone acetate treatment on homocysteine levels in women with polycystic ovary syndrome. *Arch Gynecol Obstet* 2008; **277**:25–30.
- Guzelmeric K, Alkan N, Pirimoglu M, Unal O, Turan C. Chronic inflammation and elevated homocysteine levels are associated with increased body mass index in women with polycystic ovary syndrome. *Gynecol Endocrinol* 2007; **23**:505–510.
- Hamurcu Z, Bayram F, Kahrman G, Donmez-Altuntas H, Baskol G. Micronucleus frequency in lymphocytes and 8-hydroxydeoxyguanosine level in plasma of women with polycystic ovary syndrome. *Gynecol Endocrinol* 2010; **26**:590–595.
- Harmanci A, Cinar N, Bayraktar M, Yildiz BO. Oral contraceptive plus anti-androgen therapy and cardiometabolic risk in polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2013; **78**:120–125.
- Hattersley AT, McCarthy MI. What makes a good genetic association study? *Lancet* 2005; **366**:1315–1323.
- Haydardeoglu B, Simsek E, Kilicdag EB, Bagis T. Metabolic and endocrine effects of metformin and metformin plus cyclic medroxyprogesterone acetate in women with polycystic ovary syndrome. *Int J Gynaecol Obstet* 2009; **105**:32–35.
- Hemati T, Moghadami TN, Davari TF, Salmanian B, Javadian P. High plasma homocysteine and insulin resistance in patients with polycystic ovarian syndrome. *Iran J Reprod Med* 2011; **9**:223–228.
- Heutling D, Schulz H, Nickel I, Kleinstein J, Kaltwasser P, Westphal S, Mittermayer F, Wolzt M, Krzyzanowska K, Randeja H et al. Asymmetrical dimethylarginine, inflammatory and metabolic parameters in women with polycystic ovary syndrome before and after metformin treatment. *J Clin Endocrinol Metab* 2008; **93**:82–90.
- Kadowaki T, Yamauchi T. Adiponectin and adiponectin receptors. *Endocr Rev* 2005; **26**:439–451.
- Karadeniz M, Erdogan M, Tamsel S, Zengi A, Alper GE, Caglayan O, Saygili F, Yilmaz C. Oxidative stress markers in young patients with polycystic ovary syndrome, the relationship between insulin resistances. *Exp Clin Endocrinol Diabetes* 2008; **116**:231–235.
- Karadeniz M, Erdogan M, Zengi A, Eroglu Z, Tamsel S, Olukman M, Saygili F, Yilmaz C. Methylenetetrahydrofolate reductase C677T gene polymorphism in Turkish patients with polycystic ovary syndrome. *Endocrine* 2010; **38**:127–133.
- Karadeniz M, Erdoan M, Ayhan Z, Yalcin M, Olukman M, Cetinkalp S, Alper GE, Eroglu Z, Tetik A, Cetintas V et al. Effect of G2706A and G1051A polymorphisms of the ABCA1 gene on the lipid, oxidative stress and homocysteine levels in Turkish patients with polycystic ovary syndrome. *Lipids Health Dis* 2011; **10**:1–8.
- Karaer A, Cavkaytar S, Mert I, Buyukkagnici U, Batioglu S. Cardiovascular risk factors in polycystic ovary syndrome. *J Obstet Gynaecol* 2010; **30**:387–392.
- Kaya C, Cengiz SD, Berker B, Demirtas S, Cesur M, Erdogan G. Comparative effects of atorvastatin and simvastatin on the plasma total homocysteine levels in women with polycystic ovary syndrome: a prospective randomized study. *Fertil Steril* 2009a; **92**:635–642.
- Kaya C, Erkan AF, Cengiz SD, Dunder I, Demirel OE, Bilgihan A. Advanced oxidation protein products are increased in women with polycystic ovary syndrome: relationship with traditional and nontraditional cardiovascular risk factors in patients with polycystic ovary syndrome. *Fertil Steril* 2009b; **92**:1372–1377.
- Kaya C, Cengiz SD, Satiroglu H. Obesity and insulin resistance associated with lower plasma vitamin B12 in PCOS. *Reprod Biomed Online* 2009c; **19**:721–726.
- Kaya C, Pabuccu R, Koca C, Ouz AK, Erkan AF, Korkmaz A, Erba D. Relationship between interleukin-6 levels and ambulatory blood pressure in women with polycystic ovary syndrome. *Fertil Steril* 2010a; **94**:1437–1443.
- Kaya C, Pabuccu R, Berker B, Satiroglu H. Plasma interleukin-18 levels are increased in the polycystic ovary syndrome: relationship of carotid intima-media wall thickness and cardiovascular risk factors. *Fertil Steril* 2010b; **93**:1200–1207.
- Kaya C, Akgul E, Pabuccu R. C-reactive protein and homocysteine levels are associated with abnormal heart rate recovery in women with polycystic ovary syndrome. *Fertil Steril* 2010c; **94**:230–235.
- Kaya C, Pabuccu R, Cengiz SD, Dunder I. Comparison of the effects of atorvastatin and simvastatin in women with polycystic ovary syndrome: a prospective, randomized study. *Exp Clin Endocrinol Diabetes* 2010d; **118**:161–166.
- Kazerouni T, Asadi N, Dehbashi S, Zolghadri J. Effect of folic acid in women with and without insulin resistance who have hyperhomocysteinemic polycystic ovary syndrome. *Int J Gynaecol Obstet* 2008; **101**:156–160.
- Kebapcilar L, Yuksel A, Bozkaya G, Taner CE, Kebapcilar AG, Bilgir O, Alacacioglu A, Sari I. Effects of an EE/CA compared with EE/CA-metformin on serum ADMA levels in women with polycystic ovary syndrome. *Cent Eur J Med* 2009; **4**:423–427.
- Kilic-Okman T, Kucuk M. N-acetyl-cysteine treatment for polycystic ovary syndrome. *Int J Gynecol Obstet* 2004; **85**:296–297.



- Kilic-Okman T, Guldiken S, Kucuk M. Relationship between homocysteine and insulin resistance in women with polycystic ovary syndrome. *Endocr J* 2004; **51**:505–508.
- Kilic S, Yilmaz N, Zulfikaroglu E, Erdogan G, Aydin M, Batioglu S. Inflammatory-metabolic parameters in obese and nonobese normoandrogenemic polycystic ovary syndrome during metformin and oral contraceptive treatment. *Gynecol Endocrinol* 2011; **27**:622–629.
- Kilicdag EB, Bagis T, Zeyneloglu HB, Tarim E, Aslan E, Haydardedeoglu B, Erkanli S. Homocysteine levels in women with polycystic ovary syndrome treated with metformin versus rosiglitazone: a randomized study. *Hum Reprod* 2005a; **20**:894–899.
- Kilicdag EB, Bagis T, Tarim E, Aslan E, Erkanli S, Simsek E, Haydardedeoglu B, Kucuk E. Administration of B-group vitamins reduces circulating homocysteine in polycystic ovarian syndrome patients treated with metformin: a randomized trial. *Hum Reprod* 2005b; **20**:1521–1528.
- Kirchengast S, Peterson B, Hauser G, Knogler W. Body composition characteristics are associated with the bone density of the proximal femur end in middle- and old-aged women and men. *Maturitas* 2001; **39**:133–145.
- Kisakol G, Guney E, Bayraktar F, Yilmaz C, Kabalak T, Ozmen D. Effect of surgical weight loss on free radical and antioxidant balance: a preliminary report. *Obes Surg* 2002; **12**:795–800.
- Knochenhauer ES, Key TJ, Kahsar-Miller M, Waggoner W, Boots LR, Azziz R. Prevalence of the polycystic ovary syndrome in unselected black and white women of the southeastern United States: a prospective study. *J Clin Endocrinol Metab* 1998; **83**:3078–3082.
- Konukoglu D, Serin O, Ercan M, Turhan MS. Plasma homocysteine levels in obese and non-obese subjects with or without hypertension; its relationship with oxidative stress and copper. *Clin Biochem* 2003; **36**:405–408.
- Kucuk M, Kilic-Okman T. Hormone profiles and clinical outcome after laparoscopic ovarian drilling in women with polycystic ovary syndrome. *Med Sci Monit* 2005; **11**:CR29–34.
- Kulkarni S, Wilma Delphine SCR, Raghavendra DS. Evaluation of serum C-reactive protein and oxidative stress as cardiovascular risk predictors in polycystic ovarian syndrome. *Biomedicine* 2011; **31**:225–230.
- Kurdoglu Z, Ozkol H, Tuluze Y, Koyuncu I. Oxidative status and its relation with insulin resistance in young non-obese women with polycystic ovary syndrome. *J Endocrinol Invest* 2012; **35**:317–321.
- Kucuk NK, Var A. Oxidative stress but not endothelial dysfunction exists in non-obese, young group of patients with polycystic ovary syndrome. *Acta Obstet Gynecol Scand* 2009; **88**:612–617.
- Lakhani K, Kay AR, Leiper J, Barry JA, Hardiman PJ. Symmetric dimethylarginine (SDMA) is raised in women with polycystic ovary syndrome: a pilot study. *J Obstet Gynaecol* 2011; **31**:417–419.
- Lee JY, Baw C-K, Gupta S, Aziz N, Agarwal A. Role of oxidative stress in polycystic ovary syndrome. *Curr Womens Health Rev* 2010; **6**:96–107.
- Liu S, Navarro G, Mauvais-Jarvis F. Androgen excess produces systemic oxidative stress and predisposes to beta-cell failure in female mice. *PLoS One* 2010; **5**:e11302.
- Loverro G, Lorusso F, Mei L, Depalo R, Cormio G, Selvaggi L. The plasma homocysteine levels are increased in polycystic ovary syndrome. *Gynecol Obstet Invest* 2002; **53**:157–162.
- Luque-Ramirez M, Mendieta-Azcona C, del Rey Sanchez JM, Maties M, Escobar-Morreale HF. Effects of an antiandrogenic oral contraceptive pill compared with metformin on blood coagulation tests and endothelial function in women with the polycystic ovary syndrome: influence of obesity and smoking. *Eur J Endocrinol* 2009; **160**:469–480.
- Macut D, Damjanovic S, Panidis D, Spanos N, Glisic B, Petakov M, Rousso D, Kourtis A, Bjekic J, Milic N. Oxidised low-density lipoprotein concentration—early marker of an altered lipid metabolism in young women with PCOS. *Eur J Endocrinol* 2006; **155**:131–136.
- Macut D, Simic T, Lissounov A, Pljesa-Ercegovac M, Bozic I, Djukic T, Bjekic-Macut J, Matic M, Petakov M, Suvakov S et al. Insulin resistance in non-obese women with polycystic ovary syndrome: relation to byproducts of oxidative stress. *Exp Clin Endocrinol Diabetes* 2011; **119**:451–455.
- Makedos A, Goulis DG, Papanikolaou A, Panidis D. Serum high-sensitivity C-reactive protein and homocysteine changes during hormonal therapy in women with polycystic ovary syndrome: a prospective, matched study. *Angiology* 2010; **61**:595–601.
- Mancini F, Cianciosi A, Marchesini Reggiani G, Facchinetti F, Battaglia C, de Aloysio D. Endothelial function and its relationship to leptin, homocysteine, and insulin resistance in lean and overweight eumenorrheic women and PCOS patients: a pilot study. *Fertil Steril* 2009; **91**:2537–2544.
- Mancini F, Cianciosi A, Persico N, Facchinetti F, Busacchi P, Battaglia C. Drospirenone and cardiovascular risk in lean and obese polycystic ovary syndrome patients: a pilot study. *Am J Obstet Gynecol* 2010; **202**:169.e161–169.e168.
- March WA, Moore VM, Willson KJ, Phillips DI, Norman RJ, Davies MJ. The prevalence of polycystic ovary syndrome in a community sample assessed under contrasting diagnostic criteria. *Hum Reprod* 2010; **25**:544–551.
- Markou A, Androulakis II, Mourmouris C, Tsikini A, Samara C, Sougoultzis S, Piaditis G, Kaltsas G. Hepatic steatosis in young lean insulin resistant women with polycystic ovary syndrome. *Fertil Steril* 2010; **93**:1220–1226.
- Masharani U, Gjerde C, Evans JL, Youngren JF, Goldfine ID. Effects of controlled-release alpha lipoic acid in lean, nondiabetic patients with polycystic ovary syndrome. *J Diabetes Sci Technol* 2010; **4**:359–364.
- Masharani UB, Maddux BA, Li X, Sakkas GK, Mulligan K, Schambelan M, Goldfine ID, Youngren JF. Insulin resistance in non-obese subjects is associated with activation of the JNK pathway and impaired insulin signaling in skeletal muscle. *PLoS One* 2011; **6**:e19878.
- Matsuoka T, Kajimoto Y, Watada H, Kaneto H, Kishimoto M, Umayahara Y, Fujitani Y, Kamada T, Kawamori R, Yamasaki Y. Glycation-dependent, reactive oxygen species-mediated suppression of the insulin gene promoter activity in HIT cells. *J Clin Invest* 1997; **99**:144–150.
- Michelmore KF, Balen AH, Dunger DB, Vessey MP. Polycystic ovaries and associated clinical and biochemical features in young women. *Clin Endocrinol (Oxf)* 1999; **51**:779–786.
- Mohamadin AM, Habib FA, Elahi TF. Serum paraoxonase I activity and oxidant/antioxidant status in Saudi women with polycystic ovary syndrome. *Pathophysiology* 2010a; **17**:189–196.
- Mohamadin AM, Habib FA, Al-Saggaf AA. Cardiovascular disease markers in women with polycystic ovary syndrome with emphasis on asymmetric dimethylarginine and homocysteine. *Ann Saudi Med* 2010b; **30**:278–283.
- Mohan SK, Vishnu PV. Lipid peroxidation, glutathione, ascorbic acid, vitamin E, antioxidant enzyme and serum homocysteine status in patients with polycystic ovary syndrome. *Biol Med* 2009; **1**:44–49.
- Moini A, Sarrafion F, Ziaei S, Faghihzadeh S, Arabipour A. A possible role for familial thrombophilia in women with polycystic ovarian syndrome. *Gynecol Endocrinol* 2011; **27**:80–85.
- Moini A, Tadayon S, Tehrani A, Yeganeh LM, Akhoond MR, Yazdi RS. Association of thrombophilia and polycystic ovarian syndrome in women with history of recurrent pregnancy loss. *Gynecol Endocrinol* 2012; **28**:590–593.
- Moran LJ, Hutchison SK, Meyer C, Zoungas S, Teede HJ. A comprehensive assessment of endothelial function in overweight women with and without polycystic ovary syndrome. *Clin Sci (Lond)* 2009; **116**:761–770.
- Moran LJ, Cameron JD, Strauss BJ, Teede HJ. Vascular function in the diagnostic categories of polycystic ovary syndrome. *Hum Reprod* 2011; **26**:2192–2199.
- Morgante G, La Marca A, Setacci F, Setacci C, Petraglia F, De Leo V. The cardiovascular risk factor homocysteine is not elevated in young women with hyperandrogenism or hypoestrogenism. *Gynecol Obstet Invest* 2002; **53**:200–203.
- Murri A, Garcia-Fuentes E, Garcia-Almeida JM, Garrido-Sanchez L, Mayas MD, Bernal R, Tinahones FJ. Changes in oxidative stress and insulin resistance in morbidly obese patients after bariatric surgery. *Obes Surg* 2010; **20**:363–368.
- Nacul AP, Andrade CD, Schwarz P, de Bittencourt PIH Jr, Spritzer PM. Nitric oxide and fibrinogen in polycystic ovary syndrome: associations with insulin resistance and obesity. *Eur J Obstet Gynecol Reprod Biol* 2007; **133**:191–196.
- Nafiyev Y, Sevtpat K, Muammer D, Emre O, Senol K, Leyla M. The effect of serum and intrafollicular insulin resistance parameters and homocysteine levels of nonobese, nonhyperandrogenemic polycystic ovary syndrome patients on *in vitro* fertilization outcome. *Fertil Steril* 2010; **93**:1864–1869.
- Ngo DT, Chan WP, Rajendran S, Heresztyn T, Amarasekera A, Sverdlow AL, O'Loughlin PD, Morris HA, Chirkov YY, Norman RJ et al. Determinants of insulin responsiveness in young women: impact of polycystic ovarian syndrome, nitric oxide, and vitamin D. *Nitric Oxide* 2011; **25**:326–330.
- Nguyen MT, Satoh H, Faveyukis S, Babendure JL, Imamura T, Sbdio JI, Zalevsky J, Dahiyat BI, Chi NW, Olefsky JM. JNK and tumor necrosis factor-alpha mediate free fatty acid-induced insulin resistance in 3T3-L1 adipocytes. *J Biol Chem* 2005; **280**:35361–35371.

- Oktem M, Ozcimen EE, Uckuyu A, Esinler I, Pamuk B, Bayraktar N, Kulaksizoglu S, Zeyneloglu HB. Polycystic ovary syndrome is associated with elevated plasma soluble CD40 ligand, a marker of coronary artery disease. *Fertil Steril* 2009; **91**:2545–2550.
- Olusi SO. Obesity is an independent risk factor for plasma lipid peroxidation and depletion of erythrocyte cytoprotective enzymes in humans. *Int J Obes Relat Metab Disord* 2002; **26**:1159–1164.
- Orio F Jr, Palomba S, Di Biase S, Colao A, Tauchmanova L, Savastano S, Labella D, Russo T, Zullo F, Lombardi G. Homocysteine levels and C677T polymorphism of methylenetetrahydrofolate reductase in women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2003; **88**:673–679.
- Ozata M, Mergen M, Oktenli C, Aydin A, Sanisoglu SY, Bolu E. Increased oxidative stress and hypozincemia in male obesity. *Clin Biochem* 2002; **35**:627–631.
- Ozgurtas T, Oktenli C, Dede M, Tapan S, Kenar L, Sanisoglu SY, Yesilova Z, Yenen MC, Erbil MK, Baser I. Metformin and oral contraceptive treatments reduced circulating asymmetric dimethylarginine (ADMA) levels in patients with polycystic ovary syndrome (PCOS). *Atherosclerosis* 2008; **200**:336–344.
- Palacio JR, Iborra A, Ulcova-Galova Z, Badia R, Martinez P. The presence of antibodies to oxidative modified proteins in serum from polycystic ovary syndrome patients. *Clin Exp Immunol* 2006; **144**:217–222.
- Palep-Singh M, Picton HM, Yates ZR, Barth J, Balen AH. Polycystic ovary syndrome and the single nucleotide polymorphisms of methylenetetrahydrofolate reductase: a pilot observational study. *Hum Fertil (Camb)* 2007; **10**:33–41.
- Palep-Singh M, Picton HM, Yates ZR, Barth JH, Balen AH. Plasma homocysteine concentrations and the single nucleotide polymorphisms in the methionine synthase gene (MTR 2756A>G): associations with the polycystic ovary syndrome: an observational study. *Eur J Obstet Gynecol Reprod Biol* 2008; **138**:180–186.
- Palomba S, Falbo A, Giallauria F, Russo T, Tolino A, Zullo F, Colao A, Orio F. Effects of metformin with or without supplementation with folate on homocysteine levels and vascular endothelium of women with polycystic ovary syndrome. *Diabetes Care* 2010; **33**:246–251.
- Pamuk BO, Torun AN, Kulaksizoglu M, Ertugrul D, Ciftci O, Kulaksizoglu S, Yildirim E, Demir NG. Asymmetric dimethylarginine levels and carotid intima-media thickness in obese patients with polycystic ovary syndrome and their relationship to metabolic parameters. *Fertil Steril* 2010; **93**:1227–1233.
- Pollack M, Leeuwenburgh C. Molecular mechanisms of oxidative stress in aging: free radicals, aging, antioxidants and disease. In: Sen CK, Packer L, Hanninen O (eds). *Handbook of Oxidants and Antioxidants in Exercise*. Amsterdam: Elsevier Science B.V., 1999, 881–923.
- Rajagopal G, Reddy AP, Venkata Harinarayan C, Suresh V, Bitla A, PVLN Rao S, Sachan A. Effect of lifestyle modification and metformin therapy on emerging cardiovascular risk factors in overweight Indian women with polycystic ovary syndrome. *Metab Syndr Relat Disord* 2012; **10**:273–279.
- Rajendran S, Willoughby SR, Chan WP, Liberts EA, Heresztyn T, Saha M, Marber MS, Norman RJ, Horowitz JD. Polycystic ovary syndrome is associated with severe platelet and endothelial dysfunction in both obese and lean subjects. *Atherosclerosis* 2009; **204**:509–514.
- Randeva HS, Lewandowski KC, Drzewoski J, Brooke-Wavell K, O'Callaghan C, Czupryniak L, Hillhouse EW, Prelevic GM. Exercise decreases plasma total homocysteine in overweight young women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2002; **87**:4496–4501.
- The Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome (PCOS). *Hum Reprod* 2004; **19**:41–47.
- Rudich A, Tirosch A, Potashnik R, Hemi R, Kanety H, Bashan N. Prolonged oxidative stress impairs insulin-induced GLUT4 translocation in 3T3-L1 adipocytes. *Diabetes* 1998; **47**:1562–1569.
- Rzeczynska IJ, Foyouzi N, Piotrowski PC, Celik-Ozenci C, Cress A, Duleba AJ. Antioxidants induce apoptosis of rat ovarian theca-interstitial cells. *Biol Reprod* 2011; **84**:162–166.
- Sabuncu T, Vural H, Harma M. Oxidative stress in polycystic ovary syndrome and its contribution to the risk of cardiovascular disease. *Clin Biochem* 2001; **34**:407–413.
- Sahin Y, Unluhizarci K, Yilmazsoy A, Yikilmaz A, Aygen E, Kelestimur F. The effects of metformin on metabolic and cardiovascular risk factors in nonobese women with polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2007; **67**:904–908.
- Salehpour S, Manzor-al-ajdad O, Samani EN, Abadi A. Evaluation of homocysteine levels in patients with polycystic ovarian syndrome. *Int J Fertil Steril* 2011; **4**:168–171.
- San Millan JL, Alvarez-Blasco F, Luque-Ramirez M, Botella-Carretero JL, Escobar-Morreale HF. The PON1-108C/T polymorphism, and not the polycystic ovary syndrome, is an important determinant of reduced serum paraoxonase activity in premenopausal women. *Hum Reprod* 2006; **21**:3157–3161.
- Sanchon R, Gambineri A, Alpanes M, Martinez-Garcia MA, Pasquali R, Escobar-Morreale HF. Prevalence of functional disorders of androgen excess in unselected premenopausal women: a study in blood donors. *Hum Reprod* 2012; **27**:1209–1216.
- Schachter M, Raziel A, Friedler S, Strassburger D, Bern O, Ron-El R. Insulin resistance in patients with polycystic ovary syndrome is associated with elevated plasma homocysteine. *Hum Reprod* 2003; **18**:721–727.
- Schachter M, Raziel A, Strassburger D, Rotem C, Ron-El R, Friedler S. Prospective, randomized trial of metformin and vitamins for the reduction of plasma homocysteine in insulin-resistant polycystic ovary syndrome. *Fertil Steril* 2007; **88**:227–230.
- Skrha J, Sindelka G, Kvasnicka J, Hilgertova J. Insulin action and fibrinolysis influenced by vitamin E in obese type 2 diabetes mellitus. *Diabetes Res Clin Pract* 1999; **44**:27–33.
- Soares GM, Vieira CS, Martins WP, Franceschini SA, dos Reis RM, Silva de Sa MF, Ferriani RA. Increased arterial stiffness in nonobese women with polycystic ovary syndrome (PCOS) without comorbidities: one more characteristic inherent to the syndrome? *Clin Endocrinol (Oxf)* 2009; **71**:406–411.
- Sova H, Morin-Papunen L, Puistola U, Karihtala P. Distinctly low levels of serum 8-hydroxydeoxyguanosine in women with polycystic ovary syndrome. *Fertil Steril* 2010; **94**:2670–2673.
- Soyman Z, Noyan V, Tulmac M, Yucel A, Sagsoz N, Bayrak T, Bayrak A, Cakir E. Serum paraoxonase I activity, asymmetric dimethylarginine levels, and brachial artery flow-mediated dilatation in women with polycystic ovary syndrome. *Fertil Steril* 2011; **95**:1067–1072.
- Sydow K, Munzel T. ADMA and oxidative stress. *Atheroscler Suppl* 2003; **4**:41–51.
- Takeda E, Arai H, Yamamoto H, Okumura H, Taketani Y. Control of oxidative stress and metabolic homeostasis by the suppression of postprandial hyperglycemia. *J Med Invest* 2005; **52**:259–265.
- Taskin O, Sadik S, Onoglu A, Gokdeniz R, Yilmaz I, Burak F, Wheeler JM. Adhesion formation after microlaparoscopic and laparoscopic ovarian coagulation for polycystic ovary disease. *J Am Assoc Gynecol Laparosc* 1999; **6**:159–163.
- Teede HJ, Meyer C, Hutchison SK, Zoungas S, McGrath BP, Moran LJ. Endothelial function and insulin resistance in polycystic ovary syndrome: the effects of medical therapy. *Fertil Steril* 2010; **93**:184–191.
- Temel I, Celik O, Hascalik S, Celik N, Sahin I, Aydin S. Serum nonesterified fatty acids, ghrelin, and homocysteine levels in women with polycystic ovary syndrome. *Turk J Med Sci* 2010; **40**:221–228.
- Topcu S, Caliskan M, Ozcimen EE, Tok D, Uckuyu A, Erdogan D, Gullu H, Yildirim A, Zeyneloglu H, Muderrisoglu H. Do young women with polycystic ovary syndrome show early evidence of preclinical coronary artery disease? *Hum Reprod* 2006; **21**:930–935.
- Torun AN, Vural M, Cece H, Camuzcuoglu H, Toy H, Aksoy N. Paraoxonase-I is not affected in polycystic ovary syndrome without metabolic syndrome and insulin resistance, but oxidative stress is altered. *Gynecol Endocrinol* 2011; **27**:988–992.
- Tuncman G, Hirosumi J, Solinas G, Chang L, Karin M, Hotamisligil GS. Functional *in vivo* interactions between JNK1 and JNK2 isoforms in obesity and insulin resistance. *Proc Natl Acad Sci USA* 2006; **103**:10741–10746.
- Turkcuoglu I, Engin-Ustun Y, Turan F, Kali Z, Bay Karabulut A, Meydanli M, Kafkasli A. Evaluation of asymmetric dimethylarginine, nitric oxide levels and associated independent variables in obese and lean patients with polycystic ovarian syndrome. *Gynecol Endocrinol* 2011; **27**:609–614.
- Turrens JF. Mitochondrial formation of reactive oxygen species. *J Physiol* 2003; **552**:335–344.
- Tyagi N, Sedoris KC, Steed M, Ovechkin AV, Moshal KS, Tyagi SC. Mechanisms of homocysteine-induced oxidative stress. *Am J Physiol Heart Circ Physiol* 2005; **289**:H2649–2656.
- Urakawa H, Katsuki A, Sumida Y, Gabazza EC, Murashima S, Morioka K. Oxidative stress is associated with adiposity and insulin resistance in men. *J Clin Endocrinol Metab* 2003; **88**:4673–4676.
- Uzun H, Zengin K, Taskin M, Aydin S, Simsek G, Dariyerli N. Changes in leptin, plasminogen activator factor and oxidative stress in morbidly obese patients following open and laparoscopic Swedish adjustable gastric banding. *Obes Surg* 2004; **14**:659–665.

- Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem Biol Interact* 2006; **160**:1–40.
- Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 2007; **39**:44–84.
- Van Gaal LF, Vertommen J, De Leeuw IH. The *in vitro* oxidizability of lipoprotein particles in obese and non-obese subjects. *Atherosclerosis* 1998; **137**:S39–S44.
- Verit FF, Erel O. Oxidative stress in nonobese women with polycystic ovary syndrome: correlations with endocrine and screening parameters. *Gynecol Obstet Invest* 2008; **65**:233–239.
- Verit FF, Erel O, Kocyigit A. Association of increased total antioxidant capacity and anovulation in nonobese infertile patients with clomiphene citrate-resistant polycystic ovary syndrome. *Fertil Steril* 2007; **88**:418–424.
- Victor VM, Rocha M, Banuls C, Sanchez-Serrano M, Sola E, Gomez M, Hernandez-Mijares A. Mitochondrial complex I impairment in leukocytes from polycystic ovary syndrome patients with insulin resistance. *J Clin Endocrinol Metab* 2009; **94**:3505–3512.
- Victor VM, Rocha M, Banuls C, Alvarez A, de Pablo C, Sanchez-Serrano M, Gomez M, Hernandez-Mijares A. Induction of oxidative stress and human leukocyte/endothelial cell interactions in polycystic ovary syndrome patients with insulin resistance. *J Clin Endocrinol Metab* 2011; **96**:3115–3122.
- Vincent HK, Taylor AG. Biomarkers and potential mechanisms of obesity-induced oxidant stress in humans. *Int J Obes* 2005; **30**:400–418.
- Vincent HK, Innes KE, Vincent KR. Oxidative stress and potential interventions to reduce oxidative stress in overweight and obesity. *Diabetes Obes Metab* 2007; **9**:813–839.
- Vrbikova J, Bicikova M, Tallova J, Hill M, Starka L. Homocysteine and steroids levels in metformin treated women with polycystic ovary syndrome. *Exp Clin Endocrinol Diabetes* 2002; **110**:74–76.
- Vrbikova J, Tallova J, Bicikova M, Dvorakova K, Hill M, Starka L. Plasma thiols and androgen levels in polycystic ovary syndrome. *Clin Chem Lab Med* 2003; **41**:216–221.
- Wells G, Shea B, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analysis. In: Oxford (ed.). *Proceedings of the 3rd Symposium on Systematic Reviews beyond the Basics: Improving Quality and Impact*, 2000 July 3–5; Oxford, pp. 3–5.
- Wijeyaratne CN, Nirantharakumar K, Balen AH, Barth JH, Sheriff R, Belchetz PE. Plasma homocysteine in polycystic ovary syndrome: does it correlate with insulin resistance and ethnicity? *Clin Endocrinol (Oxf)* 2004; **60**:560–567.
- Wright D, Sutherland L. Antioxidant supplementation in the treatment of skeletal muscle insulin resistance: potential mechanisms and clinical relevance. *Appl Physiol Nutr Metab* 2008; **33**:21–31.
- Yarali H, Yildirim A, Aybar F, Kabakci G, Bukulmez O, Akgul E, Oto A. Diastolic dysfunction and increased serum homocysteine concentrations may contribute to increased cardiovascular risk in patients with polycystic ovary syndrome. *Fertil Steril* 2001; **76**:511–516.
- Yildirim B, Demir S, Temur I, Erdemir R, Kaleli B. Lipid peroxidation in follicular fluid of women with polycystic ovary syndrome during assisted reproduction cycles. *J Reprod Med* 2007; **52**:722–726.
- Yildizhan R, Ilhan GA, Yildizhan B, Kulusari A, Adali E, Bugdayci G. Serum retinol-binding protein 4, leptin, and plasma asymmetric dimethylarginine levels in obese and nonobese young women with polycystic ovary syndrome. *Fertil Steril* 2011; **96**:246–250.
- Yilmaz M, Bukan N, Ersoy R, Karakoc A, Yetkin I, Ayvaz G, Cakir N, Arslan M. Glucose intolerance, insulin resistance and cardiovascular risk factors in first degree relatives of women with polycystic ovary syndrome. *Hum Reprod* 2005a; **20**:2414–2420.
- Yilmaz M, Biri A, Bukan N, Karakoc A, Sancak B, Toruner F, Pasaoglu H. Levels of lipoprotein and homocysteine in non-obese and obese patients with polycystic ovary syndrome. *Gynecol Endocrinol* 2005b; **20**:258–263.
- Yilmaz M, Bukan N, Ayvaz G, Karakoc A, Toruner F, Cakir N, Arslan M. The effects of rosiglitazone and metformin on oxidative stress and homocysteine levels in lean patients with polycystic ovary syndrome. *Hum Reprod* 2005c; **20**:3333–3340.
- Yilmaz N, Pektas M, Tonguc E, Kilic S, Gulerman C, Gungor T, Mollamahmutoglu L. The correlation of plasma homocysteine with insulin resistance in polycystic ovary syndrome. *J Obstet Gynaecol Res* 2008; **34**:384–391.
- Zawadzki JK, Dunaif A. Diagnostic criteria for polycystic ovary syndrome: towards a rational approach. In: Dunaif A, Givens JR, Haseltine FP, Merriam GR. (eds). *Polycystic Ovary Syndrome*. Boston: Blackwell Scientific Publications, 1992, 377–384.